

ENRICHED RIEMANN SPHERE, MORSE STABILITY AND EQUI-SINGULARITY IN $\mathcal{O}_2$

TZEE-CHAR KUO AND LAURENTIU PAUNESCU

ABSTRACT. The *Enriched Riemann Sphere* $\mathbb{CP}_*^1$ is $\mathbb{CP}^1$ plus a set of *infinitesimals*, having the Newton-Puiseux field $\mathbb{F}$ as coordinates. Complex Analysis is extended to the $\mathbb{F}$ -*Analysis* (*Newton-Puiseux Analysis*). The classical *Morse Stability Theorem* is also extended; the *stability idea* is used to formulate an *equi-singular deformation theorem* in $\mathbb{C}\{x, y\}(=\mathcal{O}_2)$.

A general principle we believe in is that the study of convergent power series in $n + 1$ variables is Global Analysis of polynomials in n variables.

In this paper this is illustrated in the case $n = 1$. Loosely speaking, the classical Morse Stability Theorem, *properly reformulated* in §10, and the stability notion are “transplanted” into Algebraic Curves, then applied to the classification problem of singularities.

In §1, the Riemann sphere $\mathbb{CP}^1$ is “enriched” to $\mathbb{CP}_*^1$ with “infinitesimals”, which are irreducible curve-germs, and $\mathbb{C}$ to $\mathbb{C}_*$. The Newton-Puiseux field $\mathbb{F}$ of convergent fractional power series is used as coordinates, in terms of which several structures are defined.

In §2, the Cauchy Integral Theorem, Taylor expansions, critical points, stability, *etc.*, are generalized to $\mathbb{F}$, as is the classical Morse Stability Theorem.

The notion of Morse stability for polynomials over $\mathbb{F}$ suggests a *stronger* definition for “*equi-singular deformation*” in $\mathbb{C}^2$ (Compare [25]).

For example, in contemporary Algebraic Geometry, both deformations

$$Q(x, y, t) := x^4 - t^2 x^2 y^2 + y^4, \quad P(x, y, t) := x^3 - y^4 - 3t^2 xy^{2d}, \quad d \geq 2, \quad (0.1)$$

are regarded as equi-singular: the zero sets are topologically trivial (Milnor μ -constant).

As we shall see, however, Q is *not* equi-singular from our viewpoint. The hypothesis of the Equi-singular Deformation Theorem in §3 is not satisfied. The associated family $\xi^4 - t^2 \xi^2 + 1$ is not Morse stable ($\xi = 0$ splits into three critical points when $t \neq 0$).

On the other hand, P , the Pham family ([20]), *is* equi-singular in the *sense* of Definition 2.7, although the “polar” P_x splits into $x \pm ty^d$. This is explained in Attention 8.2 and Example 2.8. The associated family $\xi^3 - 1$, being independent of t , is obviously Morse stable in the sense of Definition 5.4. Our Equi-singular Deformation Theorem applies.

Date: July 1, 2009.

2000 Mathematics Subject Classification. Primary 14HXX, 32SXX, 58K60, Secondary 58K40.

Key words and phrases. Fractional power series, Curve-germs, Infinitesimals, Newton-Puiseux analysis, Morse stability, Classification of singularities.

When does a given family $F(x, y, t)$, like Q, P above, admit a trivialization, and of what kind? An answer is given in the *Equi-singular Deformation Theorem* in §3, using the notion of Morse stability. (Similar results over $\mathbb{R}$ were announced in [15].)

The *Truncation Theorem* at the end of §3 asserts that $f(x, y)$ can be equi-singularly deformed into its “Puiseux root truncation” $\hat{f}_{root}(x, y)$. (We do not assume 0 is an isolated singularity.) This theorem is closely related to results on sufficiency of jets, like Morse Lemma, [1], [2], [5], [8], [11], [12], [18], [19], *etc.*. Compare also the classical book [9].

1. THE ENRICHED RIEMANN SPHERE

Take a holomorphic map-germ $\mathcal{A} : (\mathbb{C}, 0) \rightarrow (\mathbb{C}^2, 0)$, $\mathcal{A}(z) \neq 0$ if $z \neq 0$. The *image set-germ*, $Im(\mathcal{A})$, or the geometric locus of $\mathcal{A}$, has a well-defined tangent line $T(\mathcal{A})$ at 0. We call $Im(\mathcal{A})$ an *infinitesimal at $T(\mathcal{A}) \in \mathbb{CP}^1$* . The set of infinitesimals is denoted by $\mathbb{CP}_*^1$.

The geometric locus of $z \mapsto (az, bz)$ is identified with $[a : b] \in \mathbb{CP}^1$, hence $\mathbb{CP}^1 \subset \mathbb{CP}_*^1$.

For instance, the curve-germ $x^2 - y^3 = 0$, being the geometric locus of $z \mapsto (z^3, z^2)$, is an infinitesimal at $[0 : 1]$. It is “closer” to $[0 : 1]$ than any $[a : 1]$ is, $a \neq 0$, in the sense that its *contact order* (defined below) with $x = 0$ is higher than that between $x = ay$ and $x = 0$.

As in Projective Geometry, $\mathbb{CP}_*^1$ is the union $\mathbb{CP}_*^1 = \mathbb{C}_* \cup \mathbb{C}'_*$, where

$$\mathbb{C}_* := \{Im(\mathcal{A}) \mid T(\mathcal{A}) \neq [1 : 0]\}, \quad \mathbb{C}'_* := \{Im(\mathcal{A}) \mid T(\mathcal{A}) \neq [0 : 1]\}.$$

The classical Newton-Puiseux Theorem asserts that the field $\mathbb{F}$ of convergent fractional power series in an indeterminate y is algebraically closed. ([6], [10], [21], [22], [23].)

Recall that a non-zero element of $\mathbb{F}$ is a (finite or infinite) convergent series

$$\alpha : \alpha(y) = a_0 y^{n_0/N} + \cdots + a_i y^{n_i/N} + \cdots, \quad n_0 < n_1 < \cdots, \quad (1.1)$$

where $N \in \mathbb{Z}^+$, $n_i \in \mathbb{Z}$, $0 \neq a_i \in \mathbb{C}$. The *order* of α is $O_y(\alpha) := n_0/N$, $O_y(0) := +\infty$.

We can assume $GCD(N, n_0, n_1, \dots) = 1$. In this paper we call $m_{puis}(\alpha) := N$ the *Puiseux multiplicity* of α (for clarity). The *conjugates* of α are

$$\alpha_{conj}^{(k)}(y) := \sum a_i \theta^{kn_i} y^{n_i/N}, \quad 0 \leq k \leq N-1, \quad \theta := e^{2\pi\sqrt{-1}/N}.$$

The following $\mathbb{D}$ is an integral domain with quotient field $\mathbb{F}$ and ideals $\mathbb{M}, \mathbb{M}_1$:

$$\mathbb{D} := \{\alpha \in \mathbb{F} \mid O_y(\alpha) \geq 0\}, \quad \mathbb{M} := \{\alpha \mid O_y(\alpha) > 0\}, \quad \mathbb{M}_1 := \{\alpha \mid O_y(\alpha) \geq 1\}.$$

Define $|\alpha| := \sum 2^{-n_i/N} |a_i| (1 + |a_i|)^{-1}$. Then $d(\alpha, \beta) := |\alpha - \beta|$ is a *metric* on $\mathbb{D}$. If we fix N , then $\lim_{m \rightarrow \infty} \sum a_i(m) y^{n_i/N} = 0$ iff each $a_i(m) \rightarrow 0$ (point-wise convergence).

Given $\alpha \in \mathbb{M}_1$, let $\mathcal{A}(z) := (\alpha(z^N), z^N)$. We then define $\alpha_* := \pi_*(\alpha) := Im(\mathcal{A})$, and use $\pi_* : \mathbb{M}_1 \rightarrow \mathbb{C}_*$, a many-to-one surjective mapping, as a *coordinate system* on $\mathbb{C}_*$.

A coordinate system on $\mathbb{C}'_*$ is $\pi'_* : \mathbb{M}_1 \rightarrow \mathbb{C}'_*$, $\alpha_* := \pi'_*(\alpha) := Im(\mathcal{A})$, $\mathcal{A}(z) := (z^N, \alpha(z^N))$.

We furnish $\mathbb{C}_*$ (resp. $\mathbb{C}'_*$) with the quotient topology of π_* (resp. π'_*). As for the transition function in the overlap $\mathbb{C}_* \cap \mathbb{C}'_*$, take $x = \alpha(y)$, $n_0/N = 1$, we then “solve y in terms of x ”, obtaining $y = \beta(x) := b_0 x + b_1 x^{n'_1/N} + \cdots$, where $a_0 b_0 = 1$, each b_i is a polynomial in finitely many of $(\sqrt[n]{a_0})^{-1}$, a_1/a_0 , a_2/a_0 , ... Hence the topologies coincide in $\mathbb{C}_* \cap \mathbb{C}'_*$.

The quotient topology on $\mathbb{CP}_*^1$ is well-defined.

Let $X, Y \subset \mathbb{R}^n$ be germs of sub-analytic sets at 0, $X \cap Y = \{0\}$, $X \neq \{0\} \neq Y$. Define the *contact order* $\mathcal{C}_{ord}(X, Y)$ to be the smallest number L (the Lojasiewicz exponent) such that $d(x, y) \geq a\|(x, y)\|^L$, where $x \in X$, $y \in Y$, $\|x\| = \|y\|$, $a > 0$ constant ([4]).

In particular, for $\alpha_*, \beta_* \in \mathbb{C}P_*^1$, $\mathcal{C}_{ord}(\alpha_*, \beta_*)$ is defined; $\mathcal{C}_{ord}(\alpha_*, \alpha_*) := \infty$. Thus, in $\mathbb{C}_*$,

$$\mathcal{C}_{ord}(\alpha_*, \beta_*) = \max_j \{O_y(\alpha - \beta_{conj}^{(j)})\} = \max_{k,j} \{O_y(\alpha_{conj}^{(k)} - \beta_{conj}^{(j)})\}.$$

This is the contact order structure on $\mathbb{C}P_*^1$.

The *Puiseux (characteristic) pairs* of α ([24]), which describes the iterated torus knot of the curve-germ α_* , is denoted by $\chi_{puis}(\alpha)$ or $\chi_{puis}(\alpha_*)$.

The *Enriched Riemann Sphere* is $\mathbb{C}P_*^1$ furnished with the above structures, $\mathbb{C}_*$ is the *enriched complex plane*. (The Riemann-Zariski surface ([7], p.272) is much larger than $\mathbb{C}P_*^1$. For example, $x = y^{\sqrt{2}}$ defines a point in the former, but not in the latter.)

Convention 1.1. Throughout this paper, $\epsilon > 0$ is a sufficiently small constant,

$$I_{\mathbb{R}} := \{t \in \mathbb{R} \mid |t| < \epsilon\}, \quad I_{\mathbb{C}} := \{t \in \mathbb{C} \mid |t| < \epsilon\}, \quad I_{\mathbb{D}} := \{t \in \mathbb{D} \mid |t| < \epsilon\}.$$

We say $\varphi(w)$ is *real analytic*, $w = u + \sqrt{-1}v \in \mathbb{C}$, if it is so as a function of $(u, v) \in \mathbb{R}^2$. By “ $+\dots$ ” we mean “*plus higher order terms*”.

2. THE $\mathbb{F}$ -ANALYSIS (NEWTON-PUISEUX ANALYSIS)

Given $U \subset \mathbb{D}$, open, and $\phi : U \rightarrow \mathbb{F}$. We say ϕ is *Puiseux-Lojasiewicz bounded* if every $\alpha \in U$ has a neighbourhood $\mathcal{N}(\alpha)$ with constants $K(\alpha), L(\alpha) > 0$, such that

$$m_{puis}(\phi(\xi)) \leq K(\alpha)m_{puis}(\xi), \quad O_y(\phi(\xi)) \geq -L(\alpha), \quad \xi \in \mathcal{N}(\alpha).$$

In this paper we only study functions which are *Puiseux-Lojasiewicz bounded*.

Definition 2.1. We say ϕ is *differentiable* at $\alpha \in U$, with *derivative* $\phi'(\alpha) \in \mathbb{F}$, if

$$\phi'(\alpha) = \lim[\phi(\alpha + \delta) - \phi(\alpha)]/\delta \text{ as } \delta \rightarrow 0 \text{ } (\delta \in \mathbb{D}).$$

If $\phi'(\gamma) = 0$, γ is a *critical point*, with *multiplicity*

$$m_{crit}(\gamma) := \max\{k \mid \phi^{(i)}(\gamma) = 0, 1 \leq i \leq k\}.$$

Cauchy's Theorem. If $\phi'(\alpha)$ exists at every $\alpha \in U$, then all derivatives $\phi^{(k)}(\alpha)$ exist,

$$\oint_{z \in C} \phi(\mu) d\mu = 0, \quad \phi^{(k)}(\alpha) = \frac{k!}{2\pi\sqrt{-1}} \oint_{z \in C} \frac{\phi(\mu)}{(\mu - \alpha)^{k+1}} d\mu, \quad k \geq 0, \quad (2.1)$$

where $\mu := \alpha + z\delta$, $\delta \in \mathbb{D}$, $d\mu := \delta dz$, C a sufficiently small contour around $0 \in \mathbb{C}$.

Moreover, ϕ is $\mathbb{F}$ -analytic in the sense that if $\alpha + z\delta \in U$, $|z| < r$, $z \in \mathbb{C}$, then

$$\phi(\alpha + z\delta) = \phi(\alpha) + \dots + (1/k!)\phi^{(k)}(\alpha)(z\delta)^k + \dots, \quad |z| < r, \quad (2.2)$$

where $m_{puis}(\phi^{(k)}(\alpha)) \leq K(\alpha)$, a constant.

From now on, we consider a given $\phi : \mathbb{M}_1 \rightarrow \mathbb{M}_1$, which extends to a differentiable function $U \rightarrow \mathbb{D}$. Taking $\alpha, \delta \in \mathbb{M}_1$, $\xi := z\delta$, we have the ‘‘Taylor expansion’’ of ϕ at α :

$$\phi(\alpha + \xi) = \sum \alpha_k \xi^k, \quad \xi \in \mathbb{M}_1, \quad \alpha_k := (1/k!) \phi^{(k)}(\alpha) \in \mathbb{D}. \quad (2.3)$$

In this paper we always assume ϕ is mini-regular in ξ , say of order m , i.e.,

$$O_y(\alpha_m) = 0, \quad O_y(\alpha_k) + k \geq m \text{ for } 0 \leq k \leq m-1. \quad (2.4)$$

Thus, by the Newton-Puiseux Theorem, ϕ has m roots in $\mathbb{M}_1$,

$$Z(\phi) := \{\zeta \in \mathbb{M}_1 \mid \phi(\zeta) = 0\} := \{\zeta_1, \dots, \zeta_d\}, \quad m = \sum m_i, \quad (2.5)$$

where $m_i := m(\zeta_i)$ is the multiplicity of ζ_i . Of course ϕ has $m-1$ critical points in $\mathbb{M}_1$.

Definition 2.2. In $\mathbb{M}_1$, define $\mu \sim_\phi \nu$ iff either $\mu = \nu$ or else

$$O_y(\mu - \nu) > O_y(\mu - \zeta_i) = O_y(\nu - \zeta_i), \quad 1 \leq i \leq d.$$

The equivalence class of μ is denoted by μ_ϕ . The *height* of μ_ϕ is

$$h(\mu_\phi) := \max\{O_y(\mu - \zeta_i) \mid 1 \leq i \leq d\}.$$

The quotient space is $\mathbb{M}_{1,\phi} := \mathbb{M}_1 / \sim_\phi$, with the *contact order structure*:

$$\mathcal{C}_{ord}(\mu_\phi, \nu_\phi) := O_y(\mu - \nu) \text{ if } \mu_\phi \neq \nu_\phi; \quad \mathcal{C}_{ord}(\mu_\phi, \mu_\phi) := \infty.$$

Let $\mu_\phi(y)$ denote $\mu(y)$ with terms y^e deleted, $e > h(\mu_\phi)$; $\mu_\phi(y)$ depends only on $\mu_\phi \in \mathbb{M}_{1,\phi}$. We call $\mu_\phi(y) \in \mathbb{M}_1$ the *canonical coordinate* of $\mu_\phi \in \mathbb{M}_{1,\phi}$; μ_ϕ and $\mu_\phi(y)$ are often *identified*.

The *Puiseux pairs* of $\mu_\phi \in \mathbb{M}_{1,\phi}$ is $\chi_{puis}(\mu_\phi) := \chi_{puis}(\mu_\phi(y))$.

Example 2.3. Take $\phi(\xi) := \xi^2 - 2y^3$, $\mu(y) := y^{3/2} + y^{7/4}$. Then $h(\mu_\phi) = 3/2$, $\mu_\phi(y) = y^{3/2}$, $\chi_{puis}(\mu) = \{3/2, 7/4\}$, $\chi_{puis}(\mu_\phi) = \{3/2\}$. (We also call $\mu_\phi(y)$ the ϕ -*truncation* of $\mu(y)$.)

The *tree-model* of ϕ defined in [13] is our $\mathbb{M}_{1,\phi}$ without the structures. See §11.

Definition 2.4. In $\mathbb{M}_{1,\phi}$ define $\xi_\phi \sim_{bar} \eta_\phi$ iff either $\xi_\phi = \eta_\phi$, or else

$$h(\xi_\phi) = h(\eta_\phi) = \mathcal{C}_{ord}(\xi_\phi, \eta_\phi).$$

An equivalence class is called a *bar* (as in [13]). The *bar space* is the quotient

$$Bsp(\mathbb{M}_{1,\phi}) := \mathbb{M}_{1,\phi} / \sim_{bar}.$$

The bar containing ξ_ϕ is denoted by $B(\xi_\phi)$, having *height* $h(B(\xi_\phi)) := h(\xi_\phi)$; $B_h(\phi)$ denotes a bar of height h . The *Lojasiewicz exponent* of ϕ at ξ , or at ξ_ϕ , or on $B(\xi_\phi)$, is

$$L_\phi(\xi) := L(\xi_\phi) := L(B(\xi_\phi)) := O_y(\phi(\xi)). \quad (2.6)$$

That (2.6) is well-defined is an easy consequence of the following:

$$\phi(\xi) = \text{unit} \cdot \prod (\xi - \zeta_i)^{m_i}, \quad O_y(\phi(\xi)) = \sum m_i O_y(\xi - \zeta_i).$$

An *important special case* is $\mathcal{V} := \mathbb{M}_{1,id}$ when $\phi = id : \xi \mapsto \xi$. Here $Z(id) = \{0\}$.

We call $\mathcal{V}$ the *value space*, and $0_{\mathcal{V}} := 0_{id}$ the ‘‘zero’’ element.

If $\mu(y) = uy^h + \dots$, $u \neq 0$, then μ_{id} is *completely determined* by the pair (u, h) . Hence, if $h < \infty$, there is a *unique* bar of height h , $B_h(id) = \{(u, h) \mid u \neq 0\}$; this is a copy of $\mathbb{C} - \{0\}$.

For $h = \infty$, we have a singleton $B_\infty(id) = \{0_\nu\}$.

Definition 2.5. The *valuation function*, val , also written as val_ϕ for *clarity*, is

$$val := val_\phi : \mathbb{M}_{1,\phi} \rightarrow \mathcal{V}, \quad val(\xi_\phi) := val_\phi(\xi_\phi) := \phi(\xi)_{id}.$$

If $\phi'(\gamma) = 0$, $\gamma \in \gamma_\phi$, γ_ϕ is a *critical point* of val_ϕ . The *multiplicity*, $m_{crit}(\gamma_\phi)$, is the total number of such $\gamma \in \gamma_\phi$.

The subspace $C(val_\phi)$ of critical points is displayed as

$$C(val_\phi) := \{\gamma_{1,\phi}, \dots, \gamma_{p,\phi}\}, \quad \sum m_{crit}(\gamma_{j,\phi}) = m - 1. \quad (2.7)$$

Example. Take $\phi(\xi) = \xi^4(\xi - y)^5$, $\varepsilon \neq 0$, $h > 1$. Then $val_\phi(\varepsilon y^h + \dots) = (-\varepsilon^4, 4h + 5)$, $val_\phi((1 + \varepsilon)y + \dots) = (\varepsilon^5(1 + \varepsilon)^4, 9)$, $val_\phi(0) = val_\phi(y) = 0_\nu$.

Definition 2.6. Let ϕ be as in (2.3), (2.4). Take

$$\Phi(\xi, t) := \sum A_i(t)\xi^i \in \mathbb{F}\{\xi, t\}, \quad \Phi(\xi, 0) = \phi(\xi).$$

We call $\Phi(\xi, t) := \phi_t(\xi)$ an $\mathbb{F}$ -*analytic deformation* of ϕ .

In this paper, we always assume $A_i(t) \in \mathbb{D}\{t\}$, and Φ is *mini-regular*:

$$O_y(A_i(t)) + i \geq m, \quad 0 \leq i \leq m.$$

In $\mathbb{M}_1 \times I_\mathbb{F}$, define $(\mu, t) \sim_\Phi (\nu, t')$ iff $t = t'$, $\mu_{\phi_t} = \nu_{\phi_t}$.

The quotient space is $\mathbb{M}_1 \times_\Phi I_\mathbb{F} := \mathbb{M}_1 \times I_\mathbb{F} / \sim_\Phi$, with valuation function

$$val_\Phi : \mathbb{M}_1 \times_\Phi I_\mathbb{F} \rightarrow \mathcal{V}, \quad (\xi_{\phi_t}, t) \mapsto [\phi_t(\xi)]_{id}.$$

The subspace of critical points is

$$C(val_\Phi) := \{(\gamma_{\phi_t}, t) \in \mathbb{M}_1 \times_\Phi I_\mathbb{F} \mid \gamma_{\phi_t} \in C(val_{\phi_t})\}.$$

The *bar space* $Bsp(\mathbb{M}_1 \times_\Phi I_\mathbb{F})$ is similarly defined.

Definition 2.7. We say Φ is *almost Morse stable* if there exists a homeomorphism

$$\tau : C(val_\phi) \times I_\mathbb{F} \rightarrow C(val_\Phi), \quad (\gamma_\phi, t) \mapsto (\tau_t(\gamma_\phi), t), \quad (2.8)$$

where $L(\tau_t(\gamma_\phi)) = L(\gamma_\phi)$.

We say Φ is *Morse stable* if, in addition, the following holds.

Given $\gamma_\phi, \gamma'_\phi \in C(val_\phi)$. If $B(\gamma_\phi) = B(\gamma'_\phi)$ and $val_\phi(\gamma_\phi) = val_\phi(\gamma'_\phi)$, then

$$val_{\phi_t}(\tau_t(\gamma_\phi)) = val_{\phi_t}(\tau_t(\gamma'_\phi)), \quad B(\tau_t(\gamma_\phi)) = B(\tau_t(\gamma'_\phi)). \quad (2.9)$$

Example 2.8. For $P(x, y, t)$ in (0.1), $\Phi(\xi, t) := \xi^3 - 3t^2y^{2d}\xi - y^4$, val_{ϕ_t} has a *unique* critical point $\gamma_{\phi_t} = 0$ in $\mathbb{M}_{1,\phi_t}$, $m_{crit}(\gamma_{\phi_t}) = 2$. With $\tau_t = id$, Φ is *Morse stable*.

Attention: ϕ_t has two critical points $\xi = \pm ty^d$ in $\mathbb{M}_1$, but val_{ϕ_t} has only one in $\mathbb{M}_{1,\phi_t}$.

On the other hand, $\Psi(\xi, t) := \xi^3 + ty^3 - y^4$ is *not almost Morse stable*.

Let $\mathcal{V} \times_{Loj} Bsp(\mathbb{M}_{1,\phi}) := \{(v, B) \mid v \in \mathcal{V}, L(v) = L(B)\} \subset \mathcal{V} \times Bsp(\mathbb{M}_1, \phi)$. We may call this a “Lojasiewicz fiber product”. Define $\mathcal{V} \times_{Loj} Bsp(\mathbb{M}_1 \times_{\Phi} I_{\mathbb{F}})$ similarly.

The Morse Stability Theorem (Over $\mathbb{F}$). *Suppose Φ is Morse stable. Then there exist t -level preserving homeomorphisms (deformations) $\mathcal{D}_{\Phi}, \mathcal{D}_{\mathcal{V}}$ such that the diagram*

$$\begin{array}{ccc} \mathbb{M}_{1,\phi} \times I_{\mathbb{F}} & \xrightarrow{Val_{\phi}} & \mathcal{V} \times_{Loj} Bsp(\mathbb{M}_{1,\phi}) \times I_{\mathbb{F}} \\ \downarrow \mathcal{D}_{\Phi} & & \downarrow \mathcal{D}_{\mathcal{V}} \\ \mathbb{M}_1 \times_{\Phi} I_{\mathbb{F}} & \xrightarrow{Val_{\Phi}} & \mathcal{V} \times_{Loj} Bsp(\mathbb{M}_1 \times_{\Phi} I_{\mathbb{F}}) \end{array}$$

is commutative, where

$$Val_{\phi}(\mu_{\phi}, t) := (val_{\phi}(\mu_{\phi}), B(\mu_{\phi}), t), \quad Val_{\Phi}(\mu_{\phi_t}, t) := (val_{\phi_t}(\mu_{\phi_t}), B(\mu_{\phi_t}), t).$$

The structures are preserved. That is, if $\mathcal{D}_{\Phi}(\mu_{\phi}, t) := (\mu_{\phi,t}, t)$, $\mu_{\phi,t} \in \mathbb{M}_{1,\phi_t}$, then

$$h(\mu_{\phi,t}) = h(\mu_{\phi}), \quad \chi_{puis}(\mu_{\phi,t}) = \chi_{puis}(\mu_{\phi}), \quad \mathcal{C}_{ord}(\mu_{\phi,t}, \nu_{\phi,t}) = \mathcal{C}_{ord}(\mu_{\phi}, \nu_{\phi}); \quad (2.10)$$

and if we write $\mathcal{D}_{\mathcal{V}}(v, B, t) := (v_t, B_t, t)$, then $L(v_t) = L(v)$, $h(B_t) = h(B)$.

3. THE EQUI-SINGULAR DEFORMATION THEOREM

A real analytic map-germ $\rho : ([0, \infty), 0) \rightarrow (\mathbb{C}^2, 0)$ is called an *analytic arc* ([17]); we call the *image set-germ*, $Im(\rho)$, a *geo-arc*. The *complexification* of ρ is $\rho_{\mathbb{C}}(z) := \rho(z)$, $z \in \mathbb{C}$.

Given $k \in \mathbb{Z}^+$, define $\rho^{(k)}(s) := \rho(s^k)$. Of course, $Im(\rho^{(k)}) = Im(\rho)$, the same geo-arc.

Let $f(x, y) \in \mathbb{C}\{x, y\}$ be given, *mini-regular* in x of order m , i.e.,

$$f(x, y) = H_m(x, y) + H_{m+1}(x, y) + \cdots, \quad H_m(1, 0) \neq 0.$$

Let $\phi(\xi) := f(\xi, y)$. In $\mathbb{C}_*$, define $\alpha_* \sim_f \beta_*$ iff either $\alpha_* = \beta_*$, or else

$$\mathcal{C}_{ord}(\alpha_*, \beta_*) > \mathcal{C}_{ord}(\alpha_*, \zeta_*) = \mathcal{C}_{ord}(\beta_*, \zeta_*) \quad \forall \zeta \in Z(\phi).$$

The equivalence class of α_* is denoted by $\alpha_{*/f}$. (If $\zeta \in Z(\phi)$, $\zeta_{*/f} := \{\zeta_*\}$.)

Call $\alpha_{\phi}(y)$, and any one of the conjugates $\alpha_{\phi, conj}^{(k)}(y)$, a *canonical coordinate* of $\alpha_{*/f}$.

We say $\alpha_{*/f}, \beta_{*/f}$ are *bar equivalent*: $\alpha_{*/f} \sim_{bar} \beta_{*/f}$, if they have canonical coordinates $\alpha_{\phi}, \beta_{\phi}$ respectively, such that $\alpha_{\phi} \sim_{bar} \beta_{\phi}$.

Example 3.1. Take $f(x, y) := x^2 - 2y^3$. The curve-germs $x^2 - y^3 = 0$ and $(x^2 - y^3)^2 - xy^5 = 0$ are $\sim_f$ equivalent, hence define a same point in $\mathbb{C}_{*/f}$, with canonical coordinates $\pm y^{3/2}$.

Definition 3.2. The *valuation function* on the quotient space $\mathbb{C}_{*/f} := \mathbb{C}_*/\sim_f$ is

$$val_{*/f} : \mathbb{C}_{*/f} \rightarrow \mathcal{V}_{\mathbb{Z}}, \quad \alpha_{*/f} \mapsto \left[\prod_{k=1}^N \phi(\alpha_{\phi, conj}^{(k)}(y)) \right]_{id}, \quad N := m_{puis}(\alpha_{\phi}),$$

where $\mathcal{V}_{\mathbb{Z}} := \{0_{\mathcal{V}}\} \cup \{(u, h) \mid u \neq 0, h \in \mathbb{Z}^+\}$, a subspace of $\mathcal{V}$.

If $\phi'(\gamma) = 0, \gamma \in \mathbb{M}_1$, $\gamma_{*/f}$ is called a *critical point* of $val_{*/f}$, the *multiplicity*, $m_{crit}(\gamma_{*/f})$, is the total number of $\mu \in \mathbb{M}_1$, counting multiplicities, such that $\phi'(\mu) = 0, \mu_{*/f} = \gamma_{*/f}$.

The subspace of critical points of $val_{*/f}$ is denoted by $C(val_{*/f})$.

In $\mathbb{C}_{*/f}$, define *height* by $h(\alpha_{*/f}) := h(\alpha_\phi)$, and *contact order* by

$$\mathcal{C}_{ord}(\alpha_{*/f}, \beta_{*/f}) := \mathcal{C}_{ord}(\alpha_*, \beta_*) \text{ if } \alpha_{*/f} \neq \beta_{*/f}; \quad \mathcal{C}_{ord}(\alpha_{*/f}, \alpha_{*/f}) := \infty.$$

The *Puiseux pairs* are $\chi_{puis}(\alpha_{*/f}) := \chi_{puis}(\alpha_\phi)$.

Let $F(x, y, t) := F_t(x, y) := \sum_{i+j \geq m} c_{ij}(t)x^i y^j \in \mathbb{C}\{x, y, t\}$ be a given deformation of $f(x, y)$, i.e., $F_0(x, y) = f(x, y)$, $F_t(0, 0) \equiv 0$. In $\mathbb{C}_* \times I_{\mathbb{C}}$ define $(\alpha_*, t) \sim_F (\beta_*, t')$ iff $t = t'$ and $\alpha_{*/F_t} = \beta_{*/F_t}$. The quotient space is $\mathbb{C}_* \times_F I_{\mathbb{C}} := \mathbb{C}_* \times I_{\mathbb{C}} / \sim_F$.

The Equi-singular Deformation Theorem. *Suppose the deformation $\Phi(\xi, t) := F(\xi, y, t)$ is almost Morse stable. Then there exists a t -level preserving homeomorphism*

$$H : (\mathbb{C}^2 \times I_{\mathbb{C}}, 0 \times I_{\mathbb{C}}) \rightarrow (\mathbb{C}^2 \times I_{\mathbb{C}}, 0 \times I_{\mathbb{C}}), \quad ((x, y), t) \mapsto (H_t(x, y), t), \quad (3.1)$$

which is real bi-analytic outside $\{0\} \times I_{\mathbb{C}}$. The following hold.

- (1) $F(H_t(x, y), t) = f(x, y)$, $t \in I_{\mathbb{C}}$, i.e., $F(x, y, t)$ is “trivialized” by H .
- (2) There exists $c > 0$, $c \leq \|H_t(x, y)\| / \|(x, y)\| \leq 1/c$, $t \in I_{\mathbb{C}}$.
- (3) If $\rho(s)$ is an analytic arc, then $H_t(\rho^{(k)}(s))$ is real analytic in (s, t) (called a “geo-arc wing”) for some $k \in \mathbb{Z}^+$. In particular, H_t is geo-arc analytic in the sense that it carries geo-arcs to geo-arcs.
- (4) Take $\alpha_{*/f} \in \mathbb{C}_{*/f}$, and any ρ such that $\text{Im}(\rho_{\mathbb{C}}) \in \alpha_{*/f}$. The geo-arc $\text{Im}(H_t \circ \rho)$ is contained in a unique curve-germ, say $\delta_* \in \mathbb{C}_*$. Then $\eta_t(\alpha_{*/f}) := \delta_{*/F_t}$ is independent of the choice of ρ . Hence $\eta_t : \mathbb{C}_{*/f} \rightarrow \mathbb{C}_{*/F_t}$ is well-defined,

$$\eta_* : \mathbb{C}_{*/f} \times I_{\mathbb{C}} \rightarrow \mathbb{C}_* \times_F I_{\mathbb{C}}, \quad (\alpha_{*/f}, t) \mapsto (\eta_t(\alpha_{*/f}), t), \quad (3.2)$$

is a homeomorphism, preserving height, contact order, and Puiseux pairs.

- (5) If $\gamma_{*/f} \in C(\text{val}_{*/f})$ then $\eta_t(\gamma_{*/f}) \in C(\text{val}_{*/F_t})$, $m_{\text{crit}}(\gamma_{*/f}) = m_{\text{crit}}(\eta_t(\gamma_{*/f}))$.

Assume $\Phi(\xi, t)$ is Morse stable. Take $\gamma_{*/f}, \gamma'_{*/f} \in C(\text{val}_{*/f})$, $\gamma_{*/f} \sim_{\text{bar}} \gamma'_{*/f}$. Then

$$\text{val}_{*/f}(\gamma_{*/f}) = \text{val}_{*/f}(\gamma'_{*/f}) \text{ implies } \text{val}_{*/F_t}(\eta_t(\gamma_{*/f})) = \text{val}_{*/F_t}(\eta_t(\gamma'_{*/f})), \quad (3.3)$$

where $\eta_t(\gamma_{*/f}) \sim_{\text{bar}} \eta_t(\gamma'_{*/f})$.

Now let us consider $f(x, y)$, $Z(\phi) := \{\zeta_1, \dots, \zeta_d\}$ as in (2.5), $m_i := m(\zeta_i)$,

$$f(x, y) = u(x, y) \cdot \prod_{i=1}^d (x - \zeta_i(y))^{m_i}, \quad u(0, 0) \neq 0.$$

Let $e_i := \max_{j \neq i} \{O_y(\zeta_i - \zeta_j)\}$. Let $\hat{\zeta}_i(y)$ denote $\zeta_i(y)$ with all terms y^e deleted, $e > e_i$.

Definition 3.3. The *Puiseux root truncation* of $f(x, y)$ is, by definition,

$$\hat{f}_{\text{root}}(x, y) := \begin{cases} \prod_{i=1}^d (x - \hat{\zeta}_i(y))^{m_i} & \text{if } d > 1, \\ (x - \zeta_1)^{m_1} & \text{if } d = 1. \end{cases}$$

Let $R_i(y) := \zeta_i(y) - \hat{\zeta}_i(y)$ (the remainder). Take $u(x, y, t)$ (a deformation of unit),

$$u(0, 0, t) \neq 0, \quad u(x, y, 0) = u(x, y), \quad u(x, y, 1) = 1.$$

The *Puiseux root deformation* of $f(x, y)$ is, by definition,

$$F_{\text{root}}(x, y, t) := u(x, y, t) \cdot \prod_{i=1}^d [x - \zeta_i(y) + tR_i(y)]^{m_i} \in \mathbb{C}\{x, y, t\}.$$

Note that $\hat{f}_{\text{root}}(x, y) \in \mathbb{C}\{x, y\}$, since it is invariant under the conjugations; if $d > 1$ then $\hat{f}_{\text{root}}(x, y)$ is a polynomial. The following theorem is proved at the end of §9.

The Truncation Theorem. *The Puiseux root deformation $F_{\text{root}}(x, y, t)$ is Morse stable. In particular, $f(x, y)$ and $\hat{f}_{\text{root}}(x, y)$ are geo-arc analytically equivalent.*

Example 3.4. For a weighted expansion $f(x, y) := W_d(x, y) + \dots$, if W_d is non-degenerate then $\hat{f}_{\text{root}}(x, y) = W_d(x, y)$, $f(x, y)$ is geo-arc analytically equivalent to its initial form.

Next, $g(x, y) := (x^2 - y^4)^2 - y^{10} + \dots$ has Puiseux roots $x = \pm y^2 \pm \frac{1}{2}y^3 + \dots$,

$$\hat{g}_{\text{root}}(x, y) = (x^2 - y^4)^2 - \frac{1}{4}y^6[(x - y^2)^2 + (x + y^2)^2] + \frac{1}{16}y^{12}.$$

Note that $\hat{g}_{\text{root}}(x, y)$ is *not* obtained by deleting certain terms of $g(x, y)$.

Remark 3.5. We call $\alpha_{*/f}$ an “*f-blurred*” *infinitesimal*: Points of $\mathbb{C}_*$ equivalent under $\sim_f$ are *no longer distinguishable* – “*f-blurred*”. The notion of blurring plays a vital role in this paper. For example, the Pham deformations

$$P_{\text{even}}(x, y, t) = x^3 - y^4 + 3txy^{2d}, \quad P_{\text{odd}}(x, y, t) = x^3 - y^4 + 3txy^{2d+1},$$

where $d \geq 2$, are regarded by some experts as substantially different, since the polars are very different when $t \neq 0$. To us, however, there is only one critical point – a “*blurred polar*” – in either case, of multiplicity 2. (Compare Attention 8.2). The above theorem applies.

Remark 3.6. The above (3.3) says that the family $\{val_{*/F_t}\}$ on the spaces $\{\mathbb{C}_{*/F_t}\}$ is Morse stable, the deformation of the critical points being given by η_t . Hence, as in the classical case, we can construct a trivialization of the family, like (D_t, d_t) , in §10.

However, we are not saying that this trivialization coincides with η_* . We believe it would be too good (too strong) for this to be true.

4. PROOF OF CAUCHY’S THEOREM

Let $\delta \in \mathbb{D}$ be given, and fixed. As ϕ is Puiseux-Lojasiewicz bounded, we can write

$$y^L \phi(\alpha + z\delta) = \sum_{i=0}^{\infty} c_{\delta,i}(z) y^{n_i/K}, \quad 0 < K \leq n_0 < \dots,$$

where L, K are constants, $|z|$ sufficiently small.

Take an increment Δz and compute the derivative. We find

$$y^L \phi'(\alpha + z\delta) = \frac{1}{\delta} \sum c'_{\delta,i}(z) y^{n_i/K}. \quad (4.1)$$

In particular, $c'_{\delta,i}(z)$ exists, $c_{\delta,i}(z)$ is holomorphic,

$$c_{\delta,i}(0) = \frac{1}{2\pi\sqrt{-1}} \oint_{z \in C} \frac{c_{\delta,i}(z)}{z} dz, \quad \phi(\alpha) = \frac{1}{2\pi\sqrt{-1}} \oint_{z \in C} \frac{\phi(\mu)}{\mu - \alpha} d\mu,$$

where $\mu := \alpha + z\delta$. Then, as in Complex Analysis, (2.1) follows.

Next we show (2.2). Take $\delta = 1$ in (4.1), and then set $z = 0$. We have

$$y^L \phi'(\alpha + z) = \sum c'_{1,i}(z) y^{n_i/K}, \quad c'_{1,i}(0) = \frac{1}{\delta} c'_{\delta,i}(0).$$

Applying the same argument to higher derivatives, we have

$$c_{1,i}^{(k)}(0) = \delta^{-k} c_{\delta,i}^{(k)}(0), \quad k \geq 1.$$

Hence,

$$y^L \phi(\alpha + z\delta) = \sum_i \sum_k \frac{1}{k!} c_{\delta,i}^{(k)}(0) z^k y^{n_i/K} = \sum_k \frac{1}{k!} \phi^{(k)}(\alpha) \cdot (z\delta)^k.$$

Take $\delta = y^h$, $h \in \mathbb{Q}^+$. If $m_{\text{puis}}(\phi^{(k)}(\alpha))$ were unbounded, then $m_{\text{puis}}(\phi(\alpha + cy^h)) = \infty$ for generic c , a contradiction. This completes the proof.

Corollary 4.1. *Suppose $\phi : U \rightarrow \mathbb{D}$ is $\mathbb{F}$ -analytic, $0 \in U$, $\phi(0) = 0$. Then there exist a holomorphic germ $f(x, y)$ and $N \in \mathbb{Z}^+$ such that $\phi(\xi) = f(\xi, y^{1/N})$, $\xi \in \mathbb{M}$.*

Proof. Take $\alpha = 0$, $S := \sup_k \{m_{\text{puis}}(\phi^{(k)}(0))\}$, $N := S!$. All $\phi^{(k)}(0)(y^N)$ in (2.2) are integral power series of y , $f(x, y) := \sum (1/k!) \phi^{(k)}(0)(y^N) x^k$ is holomorphic 0, $f(0, 0) = 0$. $\square$

5. NEWTON POLYGON AT AN INFINITESIMAL

Given ϕ and α . For a term $\alpha_k \xi^k \neq 0$ in (2.3), plot a “Newton dot” at (k, q) in the (u, v) -plane, $q := O_y(\alpha_k)$. Consider the convex hull generated by $\{(a_i + u, b_i + v) \mid u, v \geq 0\}$, (a_i, b_i) the Newton dots. *The boundary $\mathcal{NP}(\phi, \alpha)$ is the Newton Polygon of ϕ at α .* (See [14].)

In the case $\phi(\xi, y) := f(\xi, y)$, $f(x, y) \in \mathbb{C}\{x, y\}$, we clearly have $\mathcal{NP}(\phi, \alpha) = \mathcal{NP}(\phi, \alpha_{\text{conj}}^{(k)})$. Hence $\mathcal{NP}(\phi, \alpha_*) := \mathcal{NP}(\phi, \alpha_{\text{conj}}^{(k)})$ is well-defined. *This is the Newton Polygon of ϕ at α_* .*

Consider $\mathcal{NP}(\phi, \alpha)$. The edges and angles are $E_i := E_i(\alpha)$, $\theta_i := \theta_{E_i}$, respectively, $\theta_{i-1} < \theta_i$, $0 \leq i \leq l$. The first edge E_0 is horizontal, the last edge E_l is vertical. Denote the right vertex of E_i by $V_i := (m_i, q_i)$, and the straight line prolonging E_i by $\mathcal{L}(E_i)$.

The vertical edge E_l is not important. We call $E_{\text{top}} := E_{l-1}$ the *top* Newton edge. The left vertex of E_{l-1} is (m_l, q_l) , which is the last, and the highest, vertex of $\mathcal{NP}(\phi, \alpha)$.

Take E_i , $i \leq l-1$. A dot $(k, q) \in E_i$ represents a term $c_k y^q \xi^k$ of (2.3). Let

$$W_{E_i}(\xi, y) := \sum c_k y^q \xi^k, \quad P_{E_i}(z) := \sum c_k z^k \in \mathbb{C}[z],$$

sum taken over $(k, q) \in E_i$. We call $P_{E_i}(z)$ the *associated polynomial* of E_i .

The *height*, or *co-slope*, of E_i is

$$h(E_i) := \tan \theta_i, \quad 0 \leq i \leq l-1; \quad h(E_l) := \tan \theta_l = \infty. \quad (5.1)$$

The *Lojasiewicz exponent* on E_i is

$$L(E_i) := q_i + m_i \tan \theta_i, \quad 0 \leq i \leq l-1; \quad L(E_l) := \infty.$$

Example 5.1. (Fig.1,2) Take $\alpha = 0$. For $\phi_1(\xi) := \xi^3 + 2y\xi^2 + y^4$, $\tan \theta_1 = 1$, $\tan \theta_{top} = 3/2$, $P_{E_1}(z) = z^3 + 2z^2$, $P_{top}(z) = 2z^2 + 1$. For $\phi_2(\xi) := \xi^3 + 2y\xi^2$, $P_{top}(z) = z^3 + 2z^2$, $E_2 = E_l$.




FIGURE 1. $\mathcal{NP}(\phi_1, 0)$




FIGURE 2. $\mathcal{NP}(\phi_2, 0)$

Notation 5.2. Suppose $\mathcal{NP}(\phi, \alpha)$, $\mathcal{NP}(\phi, \beta)$ have a common edge $E_k(\alpha) = E_k(\beta)$.

We write $E_k(\alpha) \equiv E_k(\beta)$ if they have the same Newton dots, each represents a same monomial term of ϕ . We write $\mathcal{NP}(\phi, \alpha) \equiv \mathcal{NP}(\phi, \beta)$ if this is true for every edge.

Observe that if $\alpha \notin Z(\phi)$, then $m_l = 0$, and

$$h(E_{l-1}) = \max\{O(\alpha - \zeta_i)\}, \quad \zeta_i \in Z(\phi).$$

If $\alpha \sim_\phi \beta$, then $O(\alpha - \beta) > h(E_{l-1})$, and hence $\mathcal{NP}(\phi, \alpha) \equiv \mathcal{NP}(\phi, \beta)$. Thus, $P_{E_i(\alpha_\phi)}(z)$, $\mathcal{NP}(\phi, \alpha_\phi)$, $E_i(\alpha_\phi)$, etc., are all *well-defined* (independent of the choice of $\alpha \in \alpha_\phi$).

(Similarly, $\mathcal{NP}(f, \alpha_{*/f}) := \mathcal{NP}(\phi, \alpha_\phi)$ is also well-defined.)

Theorem 5.3. Take an edge E_i of $\mathcal{NP}(\phi, \alpha_\phi)$, $1 \leq i \leq l-1$. Take a critical point c of the associated polynomial $P_{E_i}(z)$, with multiplicity $m_{crit}(c)$. Then there are exactly $m_{crit}(c)$ critical points of ϕ in $\mathbb{M}_1$, counting multiplicities, of the form

$$\mu(y) = \hat{\alpha}_\phi(y) + [cy^{\tan \theta_i} + \dots], \quad (5.2)$$

where $\hat{\alpha}_\phi$ is $\alpha_\phi(y)$ with all terms y^e (if any) deleted, $e > \tan \theta_i$.

Take $\mu_\phi \in C(val_\phi)$, $\mu_\phi \neq \alpha_\phi$. There exist a unique E_i and a unique critical point $c \neq 0$ of $P_{E_i}(z)$ such that $\mu(y)$ has the form (5.2).

According to Convention 1.1, we can also write (5.2) as

$$\mu(y) = \alpha_\phi(y) + [cy^{\tan \theta_i} + \dots].$$

Theorem 5.3 is known ([13], see also [14]). We use it several times in this paper.

The Fundamental Lemma. *Suppose Φ is almost Morse stable. Then*

$$\mathcal{NP}(\phi, \gamma_\phi) = \mathcal{NP}(\phi_t, \tau_t(\gamma_\phi)), \quad \chi_{\text{puis}}(\tau_t(\gamma_\phi)) = \chi_{\text{puis}}(\gamma_\phi), \quad (5.3)$$

where $\gamma_\phi \in C(\text{val}_\phi)$ with deformation $\tau_t(\gamma_\phi) \in C(\text{val}_{\phi_t})$, as in (2.8).

The canonical coordinate $\tau_t(\gamma_\phi)(y)$ of $\tau_t(\gamma_\phi)$ is an $\mathbb{F}$ -analytic function of $t \in I_{\mathbb{F}}$. For each E_i , $i \leq l-1$, the family $\{P_{E_i(\tau_t(\gamma_\phi))}(z)\}$ is almost Morse stable, as defined below.

Moreover, if Φ is Morse stable, then so is the family $\{P_{E_i(\tau_t(\gamma_\phi))}(z)\}$.

Definition 5.4. Given a polynomial $p(z)$ and a deformation

$$p_t(x) := a_0(t)x^n + \cdots + a_n(t) \in \mathbb{K}\{t\}[x], \quad p_0(z) = p(z), \quad a(t) \neq 0,$$

where $\mathbb{K} := \mathbb{R}, \mathbb{C}$, or $\mathbb{F}$, $t \in I_{\mathbb{K}}$.

A critical point $c_0 \in \mathbb{K}$ of $p_0(x)$ is *stable* if it admits a *continuous* deformation $c_t \in \mathbb{K}$ such that $p'_t(c_t) = 0$ and $m_{\text{crit}}(c_t) = m_{\text{crit}}(c_0)$. (The deformation is then necessarily *unique*.)

Consider the following conditions (where (3) is for Algebraic Geometry):

- (1) Every critical point of $p_0(x)$ is stable.
- (2) If c_0, c'_0 are critical points of $p_0(x)$ and $p_0(c_0) = p_0(c'_0)$, then $p_t(c_t) = p_t(c'_t)$.
- (3) If $p_0(c_0) = p'_0(c_0) = 0$, i.e., c_0 is a multiple root of $p_0(z)$, then $p_t(c_t) = 0$.

We say $\{p_t\}$ is *almost Morse stable* if (1), (3) hold, and *Morse stable* if (2) also holds.

Example 5.5. Take $\mathbb{K} = \mathbb{R}$. For $p_t(x) = x^2(x^2 + t^2) \in \mathbb{R}[x]$, 0 is a critical point of p_0 which splits into three critical points in $\mathbb{C}$, one remains in $\mathbb{R}$. Thus 0 admits a *unique continuous* deformation $c_t \equiv 0$ in $\mathbb{R}$. But $m_{\text{crit}}(c_t)$ is not constant, 0 is *unstable*.

Proof. We use the “edging forward argument” to prove (5.3). The Tschirnhausen transformation is applied recursively along the edges of $\mathcal{NP}(\phi_t, \gamma_\phi)$ (“edging forward”) in order to “clear” all dots of ϕ_t lying below $\mathcal{NP}(\phi, \gamma_\phi)$. If $m_l > 0$, all dots to the left of the vertical edge E_l are also cleared. The details are as follows.

Consider $\mathcal{NP}(\phi, \gamma_\phi)$. The edges are denoted by E_i . The right vertex of E_i is (m_i, q_i) .

Let us first compare $\mathcal{NP}(\phi, \gamma_\phi)$ with $\mathcal{NP}(\phi_t, \gamma_\phi)$. Write

$$\phi_t(\gamma_\phi + \xi) = \phi(\gamma_\phi + \xi) + P_t(\gamma_\phi + \xi), \quad P_0(\gamma_\phi + \xi) \equiv 0.$$

The (non-zero) terms of P_t are represented by dots. Some may lie below $\mathcal{NP}(\phi, \gamma_\phi)$.

Suppose we already know that $P_t(\gamma_\phi + \xi)$ has no dot below the lines $\mathcal{L}(E_j)$, $0 \leq j \leq k-1$. We can then clear the dots under the line $\mathcal{L}(E_k)$ as follows.

The left vertex of E_{k-1} represents a term $ay^{q_k}\xi^{m_k}$ of $\phi(\gamma_\phi + \xi)$, which, with a Tschirnhausen transformation, can “swallow” all dots of P_t of the form $(m_k - 1, q)$, $q \in \mathbb{Q}^+$.

This means the following. There exists $\beta_t(y) \in \mathbb{M}_1$, $\mathbb{F}$ -analytic in t , such that

- (1) $O_y(\beta_t(y)) \geq \tan \theta_{k-1}$, $\beta_0(y) = 0$;
- (2) The coefficient of ξ^{m_k-1} in the Taylor expansion of $\phi(\gamma_\phi + \beta_t + \xi)$ is *independent* of t .

Indeed, $\gamma_\phi + \xi \mapsto \gamma_\phi + \xi + \beta_t$ is the *unique translation* (Tschirnhausen transformation) which has the above two properties. (Attention: No dot of $\phi(\gamma_\phi + \xi)$ has been swallowed in the process. In this way, we have $\beta_0 = 0$. This property is important.)

Now, consider $\mathcal{NP}(\phi_t, \beta_t + \gamma_\phi)$. Because of (1), $E_j(\gamma_\phi) \equiv E_j(\gamma_\phi + \beta_t)$, $j \leq k-2$, (we use Notation 5.2,) and P_t still has no dot below the line $\mathcal{L}(E_j)$, $j \leq k-1$.

Let $E'_{k-1}, E'_k, \dots$, denote the remaining edges of $\mathcal{NP}(\phi_t, \beta_t + \gamma_\phi)$. By (1), $\theta_{E_{k-1}} = \theta_{E'_{k-1}}$.

Next we show $E_{k-1} = E'_{k-1}$. Let $V'_j := (m'_j, q'_j)$ denote the right vertex of E'_j .

Suppose $E'_{k-1} \neq E_{k-1}$. Then $m'_k \leq m_k - 2$. We shall derive a contradiction.

For generic $t \in \mathbb{C}$, 0 is a root of $P_{E'_{k-1}}(z)$ of multiplicity m'_k ($m'_k \geq 0$), but when $t = 0$, the multiplicity is m_k . Hence, by an elementary argument, there exists $a(t)$ such that

$$\frac{d}{dz} P_{E'_{k-1}}(a(t)) = 0, \quad P_{E'_{k-1}}(a(t)) \neq 0, \quad \lim_{t \rightarrow 0} a(t) = 0.$$

Thus, by Theorem 5.3, ϕ_t has a critical point of the form

$$\Gamma_t(y) := [\beta_t(y) + \gamma_\phi(y)] + [a(t)y^e + \dots], \quad e := \tan \theta_{E_{k-1}}.$$

Hence $h(\Gamma_{t, \phi_t}) = \tan \theta_{E_{k-1}}$, $L_{\phi_t}(\Gamma_{t, \phi_t}) = q_k + m_k \tan \theta_{E_{k-1}}$, both are *independent* of t .

Take $\mu_\phi \in C(\text{val}_\phi)$, $\mu_\phi \neq \gamma_\phi$. Then take E_i , $c \neq 0$, for μ_ϕ as in (5.2) with γ_ϕ replacing α_ϕ . We say μ_ϕ is of the *lower kind* if $i \leq k-1$, and of the *higher kind* if $i \geq k$.

If $\mu_\phi = \gamma_\phi$, we say μ_ϕ is of the *higher kind*.

Take ε , sufficiently small. The ε -neighborhood $\mathcal{N}_\varepsilon(\gamma_\phi)$ of γ_ϕ clearly does not contain any μ_ϕ of the lower kind. On the other hand, if $|t|$ is sufficiently small, then, by continuity, $\Gamma_{t, \phi_t} \in \mathcal{N}_\varepsilon(\gamma_\phi)$. Hence $\Gamma_{t, \phi_t} \neq \tau_t(\mu_\phi)$ for any μ_ϕ of the lower kind.

If μ_ϕ is of the higher kind, then $L(\mu_\phi) = L(\tau_t(\mu_\phi)) > L(\Gamma_{t, \phi_t})$. Hence $\Gamma_{t, \phi_t} \neq \tau_t(\mu_\phi)$ for any μ_ϕ of the higher kind either. Thus we must have $E'_{k-1} = E_{k-1}$.

Next we show $\theta_{E'_k} = \theta_{E_k}$. Suppose $\theta_{E'_k} < \theta_{E_k}$. Then $m'_{k+1} \leq m_k - 2$, and there would exist $a(t)$ as above. Using the same argument we again arrive at a contradiction.

Hence the Tschirnhausen transformation $\xi \mapsto \xi + \beta_t$ clears all dots of P_t below $\mathcal{L}(E_k)$.

A recursive application of the Tschirnhausen transformations, beginning with $k = 1$, clears all dots of P_t below $\mathcal{NP}(\phi, \gamma_\phi)$.

Let $\xi \mapsto \xi + B_t$ denote their composition. We then compare the polygons:

$$\mathcal{NP}^{(0)} := \mathcal{NP}(\phi, \gamma_\phi), \quad \mathcal{NP}^{(1)} := \mathcal{NP}(\phi_t, \gamma_\phi + B_t), \quad \mathcal{NP}^{(2)} := \mathcal{NP}(\phi_t, \tau_t(\gamma_\phi)).$$

We have just proved $\mathcal{NP}^{(0)} = \mathcal{NP}^{(1)}$. Next we show $\mathcal{NP}^{(1)} = \mathcal{NP}^{(2)}$.

We can assume $B_t = 0$. This can be achieved by the substitution $\xi \rightarrow \xi + B_t(y)$.

Let us write $P_{top}^{(1)}(z) := P_{top}^{(0)}(z) + Q_t(z)$, $Q_0(z) \equiv 0$, where 0 is a critical point of $P_{top}^{(0)}(z)$.

As t varies away from 0, this critical point cannot split into two or more critical points of $P_{top}^{(1)}(z)$. For if it did, the homeomorphism τ_t cannot exist.

Hence 0 admits a *unique continuous* deformation c_t , $c_0 = 0$, which is a critical point of $P_{top}^{(1)}(z)$, $m_{crit}(c_t) = m_{puis}(c_0)$. It follows that c_t is a *simple root* of the equation

$$\frac{d^m}{dz^m} P_{top}^{(1)}(z) = 0, \quad m := m_{crit}(c_0),$$

and

$$\tau_t(\gamma_\phi)(y) = \gamma_\phi(y) + c_t y^e, \quad e := \tan \theta_{top}^{(1)}.$$

It follows that $c_t, \tau_t(\gamma_\phi)(y)$ are $\mathbb{F}$ -analytic. (The Implicit Function Theorem holds in $\mathbb{F}$.) Let the Taylor expansion of ϕ_t at γ_ϕ be $\sum_{k,q} c_{kq}(t)y^q\xi^k$. Then that at $\tau_t(\gamma_\phi)$ is

$$\sum c_{kq}(t)y^q[\xi + c_t y^e]^k = \sum c_{kq}(t)[y^q\xi^k + \dots], \quad c_0 = 0. \quad (5.4)$$

Consider $E_i^{(1)}, i \leq l-2$. Since $e > \tan \theta_i$, the terms in “ $+\dots$ ” are represented by dots lying *strictly* above all $E_i^{(1)}$. Hence $E_i^{(1)} \equiv E_i^{(2)}, i \leq l-2$. (See Notation 5.2.)

Now we show $E_{top}^{(1)} = E_{top}^{(2)}$. (But not $E_{top}^{(1)} \equiv E_{top}^{(2)}$.)

First suppose $P_{top}^{(0)}(0) \neq 0$. The left vertex of $E_{top}^{(1)} = E_{top}^{(2)}$ lies on the vertical coordinate axis. Hence $P_{top}^{(1)}(0) \neq 0, P_{top}^{(1)}(c_t) \neq 0$ ($|t|$ small). But $P_{top}^{(2)}(0) = P_{top}^{(1)}(c_t)$, hence $E_{top}^{(1)} = E_{top}^{(2)}$.

Suppose $P_{top}^{(0)}(0) = 0$, i.e., 0 is a multiple root. As $\mathcal{NP}^{(0)} = \mathcal{NP}^{(1)}$, we must have

$$c_t \equiv 0, \quad m_{crit}(c_t) = m_l = m_{puis}(\tau_t(\gamma_\phi)).$$

Hence $E_{top}^{(1)} = E_{top}^{(2)}$, and $\mathcal{NP}^{(1)} = \mathcal{NP}^{(2)}$.

It is easy to see that $\chi_{puis}(\alpha_\phi)$ can be expressed in terms of the co-slopes of the edges of $\mathcal{NP}(\phi, \alpha_\phi)$. It follows that $\chi_{puis}(\tau_t(\gamma_\phi)) = \chi_{puis}(\gamma_\phi)$. This completes the proof of (5.3).

Now assume Φ is Morse stable. We show $\{P_{E_i(\tau_t(\gamma_\phi))}(z)\}$ is Morse stable.

Take a critical point c of $P_{E_i(\gamma_\phi)}(z)$. Take μ as in (5.2). Consider $\mu_\phi, \mathcal{NP}(\phi, \mu_\phi)$, etc..

Note that $P_{E_i(\gamma_\phi)}(z+c) = P_{E_i(\mu_\phi)}(z)$ (differ merely by a translation).

First, if $P_{E_i(\gamma_\phi)}(c) \neq 0$, then $P_{E_i(\mu_\phi)}(0) \neq 0$. Hence $E_i(\mu_\phi)$ has its left vertex on the vertical axis, 0 being a critical point of $P_{E_i(\mu_\phi)}(z)$. Then, as in the argument for γ_ϕ , 0 admits a *unique continuous* deformation which is a critical point of $P_{E_i(\tau_t(\mu_\phi))}(z)$ with constant multiplicity. This says that c is a stable critical point of $P_{E_i(\tau_t(\gamma_\phi))}(z)$.

Suppose $P_{E_i(\gamma_\phi)}(c) = 0$, say of multiplicity k . Then 0 is a root of $P_{E_i(\mu_\phi)}(z)$, also of multiplicity k . Since $\mathcal{NP}(\phi, \mu_\phi) = \mathcal{NP}(\phi_t, \tau_t(\mu_\phi))$, 0 is obviously stable. Hence so is c .

Now we show (2) in Definition 5.4. Let us write $p_t(z) := P_{E_i(\tau_t(\gamma_\phi))}(z)$. Let $c \neq c'$ be critical points of $p_0(z)$. Take μ, μ' for E_i, c and c' respectively, $\mu_\phi, \mu'_\phi \in C(val_\phi)$.

First, suppose $p_0(c) = p_0(c') \neq 0$. In this case,

$$h(\mu_\phi) = h(\mu'_\phi) = O(\mu_\phi - \mu'_\phi) = \tan \theta_{E_i}.$$

Hence $B(\mu_\phi) = B(\mu'_\phi)$, $val_\phi(\mu_\phi) = val_\phi(\mu'_\phi)$. Then, by (2.9), $val_{\phi_t}(\tau_t(\mu_\phi)) = val_{\phi_t}(\tau_t(\mu'_\phi))$. That is, $(p_t(c_t), \tan \theta_{E_i}) = (p_t(c'_t), \tan \theta_{E_i}) \in \mathcal{V}$. In particular, $p_t(c_t) = p_t(c'_t)$.

Suppose $p_0(c) = p_0(c') = 0$. Then c, c' are multiple roots of p_0 , and, as shown before, their deformations remain multiple roots of p_t , $p_t(c_t) = p_t(c'_t) = 0$.

By the same argument, if Φ is almost Morse stable then so are the families $\{P_{E_i(\tau_t(\gamma_\phi))}\}$. $\square$

Corollary 5.6. Let $Z(\Phi) := \{(\zeta_t, t) \mid \zeta_t \in Z(\phi_t)\}$. There exists a bijection

$$\mathcal{D} : Z(\phi) \times I_{\mathbb{F}} \rightarrow Z(\Phi), \quad (\zeta, t) \mapsto (\zeta_t, t), \quad \zeta_0 = \zeta,$$

where $t \mapsto \zeta_t$ is $\mathbb{F}$ -analytic; the Newton Polygon $\mathcal{NP}(\phi_t, \zeta_t)$ is independent of t .

Take $\zeta, \zeta' \in Z(\phi)$, $\gamma_\phi \in C(val_\phi)$. Then $O_y(\zeta_t - \zeta'_t), O_y(\zeta_t - \tau_t(\gamma_\phi))$ are independent of t .

Proof. If ζ is a multiple root, $\zeta = \gamma_\phi$, then $\zeta_t = \tau_t(\gamma_\phi)$ is the deformation.

Otherwise, we choose $\gamma_\phi \in C(val_\phi)$ such that

$$O(\zeta - \gamma_\phi) \geq O(\zeta - \mu_\phi), \quad \forall \mu_\phi \in C(val_\phi). \quad (5.5)$$

We then have $h(\gamma_\phi) = O(\zeta - \gamma_\phi)$, and

$$\zeta(y) = \gamma_\phi(y) + [by^{h(\gamma_\phi)} + \dots], \quad P_{E(\gamma_\phi)}(b) = 0 \neq P'_{E(\gamma_\phi)}(b).$$

(If $P'_{E(\gamma_\phi)}(b) = 0$, then there would exist μ_ϕ which fails (5.5).) Using the Implicit Function Theorem we can find ζ_t , which is $\mathbb{F}$ -analytic, and $O(\zeta_t - \tau_t(\gamma_\phi))$ is constant.

Given ζ' . Take γ'_ϕ as in (5.5). Then $O(\tau_t(\gamma_\phi) - \tau_t(\gamma'_\phi))$ is constant, so is $O(\zeta_t - \zeta'_t)$. $\square$

6. RELATIONS BETWEEN BARS AND EDGES

Take a bar B , $h(B) < \infty$. Take $\beta \in \beta_\phi \in B$. Define $\zeta_B(y)$ to be $\beta(y)$ with all terms y^e deleted, $e \geq h(B)$. Clearly, $\zeta_B(y)$ depends *only* on B , not on the choices of β , β_ϕ .

Take an indeterminate z , and write

$$\phi(\zeta_B(y) + zy^{h(B)}, y) := P_B(z)y^{L(B)} + \dots, \quad P_B(z) \neq 0, \quad (6.1)$$

where $L(B)$ was defined in (2.6). We call $P_B(z)$ the associated polynomial of B .

Let $Z(P_B)$ denote the zero set of $P_B(z)$. Using the canonical coordinates, we can identify B with $\mathbb{C} - Z(P_B)$. Hence $\bar{B}$, the metric space completion of B , is a copy of $\mathbb{C}$; and

$$val_\phi(\zeta_B(y) + zy^{h(B)}) = (P_B(z), L(B)) \in \mathcal{V}, \quad z \notin Z(P_B).$$

Take $\alpha \in \mathbb{M}_1$. If $\alpha(y) = \zeta_B(y) + ay^{h(B)} + \dots$, $a \in \mathbb{C}$, we say $\bar{B}$ is a support of α ; a is the $\bar{B}$ -coordinate of α , and also of α_ϕ . Observe that $\alpha_\phi \in B$ iff $a \notin Z(P_B)$.

Notation 6.1. Write $\alpha \perp \bar{B}$ if α is supported by $\bar{B}$; $Supp(\alpha) := \{\bar{B} \mid \alpha \perp \bar{B}\}$.

Let α be given. We now define $\mathcal{NP}_{ext}(\phi, \alpha)$ by adding “vertex edges” to $\mathcal{NP}(\phi, \alpha)$.

Take a vertex $V_i = (m_i, q_i)$ of $\mathcal{NP}(\phi, \alpha)$, $m_i \geq 1$, representing a term $cy^{q_i}\xi^{m_i}$ in (2.3), where $c \neq 0$. Take $h \in \mathbb{Q}^+$, $\tan \theta_{i-1} < h < \tan \theta_i$. Let

$$E(h) := (V_i, h), \quad P_{E(h)}(z) := cz^{m_i}. \quad (6.2)$$

We call $E(h)$ a *vertex edge* and $P_{E(h)}(z)$ the *associated polynomial*. The *height*, or *co-slope*, of $E(h)$ is, by definition, $h(E(h)) := h$.

Let $\mathcal{NP}_{ext}(\phi, \alpha)$ denote the edges $\{E_0, \dots, E_{l-1}\}$ of $\mathcal{NP}(\phi, \alpha)$ plus the vertex edges.

Convention 6.2. For an edge E of $\mathcal{NP}(\phi, \alpha)$, $h := h(E) < \infty$, we also write E as $E(h)$.

Now we define

$$\iota : Supp(\alpha) \rightarrow \mathcal{NP}_{ext}(\phi, \alpha), \quad \bar{B} \mapsto \iota(\bar{B}),$$

where $\iota(\bar{B})$ is the *unique* edge of height (co-slope) $h(\iota(\bar{B})) = h(B)$. This is a bijection.

Take $\bar{B} \in Supp(\alpha)$, $h := h(B)$. Let c be the largest constant such that $\mathcal{NP}(\phi, \alpha)$ is bounded below by the line $\mathcal{L}(h) : u + v/h = c$. Let i be the smallest integer such that

$h \leq \tan \theta_{E_i}$. If $h = \tan \theta_{E_i}$, all dots on E_i lie on $\mathcal{L}(h)$. If $h < \tan \theta_{E_i}$, V_i is the *only* vertex lying on $\mathcal{L}(h)$. In either case, we define $\iota(\bar{B}) := E(h)$. See Fig.4, §11.

The corresponding associated polynomials differ merely by a translation:

$$P_B(z) = P_{\iota(\bar{B})}(z - a), \quad a \text{ the } \bar{B}\text{-coordinate of } \alpha. \quad (6.3)$$

7. PROOF OF THE MORSE STABILITY THEOREM OVER $\mathbb{F}$

Give B , $h(B) < \infty$. Take $\zeta \in Z(\phi)$, $\zeta \perp \bar{B}$. Let ζ_t be the deformation of ζ in Corollary 5.6. Define B_t to be the *unique* bar such that $h(B_t) = h(B)$, $\zeta_t \perp \bar{B}_t$.

If $\zeta' \in Z(\phi)$ and $\zeta' \perp \bar{B}$, then $O_y(\zeta_t - \zeta'_t) = O_y(\zeta - \zeta') \geq h(B)$. Hence $\text{Supp}(\zeta_t)$, $\text{Supp}(\zeta'_t)$ have the same set of bars of height $\leq h(B)$.

It follows that B_t is well-defined (*independent* of the choice of ζ),

$$\mathcal{D}_{\text{bar}} : Bsp(\mathbb{M}_{1,\phi}) \times I_{\mathbb{F}} \rightarrow Bsp(\mathbb{M}_1 \times_{\Phi} I_{\mathbb{F}}), \quad (B, t) \mapsto (B_t, t),$$

is a homeomorphism, $h(B_t) = h(B)$.

Lemma 7.1. *The family $\{P_{B_t}(z)\}$ is Morse stable (in the sense of Definition 5.4).*

Proof. In $\mathcal{NP}(\phi, \zeta)$, the left vertex of E_{top} is (m_l, q_l) , $m_l \geq 1$, where m_l is the multiplicity of ζ (as a root of ϕ). Therefore $P_{\text{top}}(0) = 0$, $\deg P_{\text{top}}(z) \geq 2$, $P_{\text{top}}(z)$ is *not* a monomial.

Hence there exists c , $P'_{\text{top}}(c) = 0 \neq P_{\text{top}}(c)$. By Theorem 5.3, there exists $\gamma_{\phi} \in C(\text{val}_{\phi})$, whose $\bar{B}$ -coordinate is c , $B := \iota^{-1}(E_{\text{top}})$. Then c has a deformation c_t , $c_0 = c$,

$$\tau_t(\gamma_{\phi})(y) = \zeta_B(y) + c_t y^{h(B)}, \quad \mathcal{NP}(\phi, \gamma_{\phi}) = \mathcal{NP}(\phi_t, \tau_t(\gamma_{\phi})).$$

Let us compare the edge $E_i(\zeta_t)$ of $\mathcal{NP}(\phi_t, \zeta_t)$ with $E_i(\tau_t(\gamma_{\phi}))$ of $\mathcal{NP}(\phi_t, \tau_t(\gamma_{\phi}))$.

If $i \leq l - 2$, then we clearly have $E_i(\zeta_t) \equiv E_i(\tau_t(\gamma_{\phi}))$. Hence $P_{E_i(\zeta_t)}(z) = P_{E_i(\tau_t(\gamma_{\phi}))}(z)$.

By the Fundamental Lemma, $\{P_{E_i(\tau_t(\gamma_{\phi}))}(z)\}$ is Morse stable. Hence if $\bar{B}' \in \text{Supp}(\zeta)$ and $h(B') < \tan \theta_{\text{top}}$, then $\{P_{B'_t}(z)\}$ is Morse stable.

As for the top edges $E_{\text{top}}(\zeta_t)$, $E_{\text{top}}(\tau_t(\gamma_{\phi}))$, their associated polynomials differ merely by a translation $z \mapsto z + c_t - a_t$, where a_t is the $\bar{B}$ -coordinate of ζ_t . The stability of the latter implies that of the former.

Finally, if $h(B'') > \tan \theta_{\text{top}}$, then $P_{B''_t}(z)$ is a monomial, hence Morse stable. $\square$

Take $B \in Bsp(\mathbb{M}_{1,\phi})$, and deformation B_t . Recall that $\bar{B} = \bar{B}_t = \mathbb{C}$. Applying the classical Morse Stability Theorem (§10) to $\{P_{B_t}\}$, we have homeomorphisms D_t , d_t such that

$$\begin{array}{ccc} \bar{B} & \xrightarrow{P_B} & \mathbb{C} \\ \downarrow D_t & & \downarrow d_t \\ \bar{B}_t & \xrightarrow{P_{B_t}} & \mathbb{C} \end{array}$$

is commutative, where D_t preserves the critical points and zeros ($d_t(0) = 0$).

Given $\alpha_{\phi} \in B$, with $\bar{B}$ -coordinate a . Take $\alpha_{\phi,t} \in B_t$ whose $\bar{B}_t$ -coordinate is $D_t(a)$:

$$\alpha_{\phi}(y) = \zeta_B(y) + ay^{h(B)}, \quad \alpha_{\phi,t}(y) = \zeta_{B_t}(y) + D_t(a)y^{h(B)}.$$

Thus, we have a homeomorphism:

$$\mathcal{D}_\Phi : \mathbb{M}_{1,\phi} \times I_\mathbb{F} \rightarrow \mathbb{M}_1 \times_\Phi I_\mathbb{F}, \quad (\alpha_\phi, t) \mapsto (\alpha_{\phi,t}, t).$$

Next, take $(u, L) \in \mathcal{V}$, $B \in Bsp(\mathbb{M}_{1,\phi})$, $L = L(B) (< \infty)$. We define

$$\mathcal{D}_\mathcal{V}((u, L), B, t) := ((d_t(u), L), B_t, t).$$

If $h(B) = \infty$, then $B = \{\zeta\}$, $\zeta \in Z(\phi)$. We define

$$\mathcal{D}_\mathcal{V}(0_\mathcal{V}, \{\zeta\}, t) := (0_\mathcal{V}, \{\zeta_t\}, t).$$

We then have $\mathcal{D}_\mathcal{V} \circ Val_\phi = Val_\Phi \circ \mathcal{D}_\Phi$.

It remains to show that $\mathcal{D}_\Phi$, $\mathcal{D}_\mathcal{V}$ preserve the structures.

Take μ_ϕ . Take $\zeta \in Z(\phi)$ such that $h(\mu_\phi) = O(\mu_\phi - \zeta)$. Then

$$h(\mu_\phi) = O(\mu_\phi - \zeta) = O(\mu_{\phi,t} - \zeta_t) = h(\mu_{\phi,t}).$$

Next, $\chi_{pvis}(\mu_\phi)$ and $\chi_{pvis}(\mu_{\phi,t})$ can be expressed in terms of the co-slopes of the edges of $\mathcal{NP}(\phi, \zeta) = \mathcal{NP}(\phi_t, \zeta_t)$. Hence $\chi_{pvis}(\mu_\phi) = \chi_{pvis}(\mu_{\phi,t})$.

As for the contact order, first suppose $\mu_\phi \sim_{bar} \nu_\phi$. Then there exists $\zeta \in Z(\phi)$,

$$h(\mu_\phi) = h(\nu_\phi) = O(\mu_\phi - \zeta) = O(\nu_\phi - \zeta),$$

which remain valid when the parameter t is added. Hence $\mathcal{C}_{ord}(\mu_\phi, \nu_\phi) = \mathcal{C}_{ord}(\mu_{\phi,t}, \nu_{\phi,t})$.

Now suppose $B(\mu_\phi) \neq B(\nu_\phi)$. Take $\zeta \perp \bar{B}(\mu_\phi)$, $\zeta' \perp \bar{B}(\nu_\phi)$. Then

$$\mathcal{C}_{ord}(\mu_\phi, \nu_\phi) = O(\zeta - \zeta') = O(\zeta_t - \zeta'_t) = \mathcal{C}_{ord}(\mu_{\phi,t}, \nu_{\phi,t}).$$

8. THE TRIVIALIZATION VECTOR FIELD

When a coordinate system $(z_1, \dots, z_n)$ of $\mathbb{C}^n$ is chosen, we use $\{\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n}\}$ to denote the standard orthonormal basis, with hermitian product

$$\langle \sum a_i \frac{\partial}{\partial z_i}, \sum b_i \frac{\partial}{\partial z_i} \rangle = \sum a_i \bar{b}_i \quad (\bar{b}_i \text{ the complex conjugate of } b_i).$$

For a holomorphic function $h(z_1, \dots, z_n)$, the *gradient* of h ([17], p.33) is

$$Grad h := \sum \frac{\partial \bar{h}}{\partial z_i} \frac{\partial}{\partial z_i}.$$

Let $f(x, y)$, $F(x, y, t)$, ϕ_t , Φ , be as in §3. To prove the Equi-singular Deformation Theorem, we use a vector field $\vec{\mathcal{F}}(x, y, t)$ which is defined in two steps, following Ehresmann's idea ([16]), where $(x, y, t) \in \mathcal{N}$, $\mathcal{N}$ a sufficiently small neighborhood of $\{0\} \times I_\mathbb{C}$ in $\mathbb{C}^2 \times I_\mathbb{C}$.

Step One. Take $\gamma_\phi \in C(val_\phi)$, with deformation $\gamma(y, t) := \tau_t(\gamma_\phi)(y)$ as in (2.8). In this step we assume $\gamma(y, t)$ is a holomorphic function in (y, t) , $\gamma(0, t) \equiv 0$.

The curve-germ defined by $x = \gamma(y, t)$ is smooth, $t \in I_\mathbb{C}$.

We define $\vec{\mathcal{F}}_\gamma(x, y, t)$ as follows. The coordinate transformation

$$\mathcal{D} : (x, y, t) \mapsto (x_\gamma, y_\gamma, t_\gamma) := (x - \gamma(y, t), y, t)$$

is holomorphic, $\mathcal{D}^{-1}$ transforms $F(x, y, t)$ to

$$F^{(\gamma)}(x_\gamma, y_\gamma, t_\gamma) := F(x_\gamma + \gamma(y_\gamma, t_\gamma), y_\gamma, t_\gamma). \quad (8.1)$$

Convention 8.1. We shall use F_{x_γ} , F_{y_γ} , F_{t_γ} to denote the partial derivatives of $F^{(\gamma)}$. The notations are simpler, but cause no confusion.

Now, consider the vector field

$$\vec{V}(x_\gamma, y_\gamma, t_\gamma) := -A(x_\gamma, y_\gamma, t_\gamma)x_\gamma \frac{\partial}{\partial x_\gamma} - B(x_\gamma, y_\gamma, t_\gamma)y_\gamma \frac{\partial}{\partial y_\gamma} + \frac{\partial}{\partial t_\gamma},$$

where

$$A := \frac{\bar{x}_\gamma \bar{F}_{x_\gamma} F_{t_\gamma}}{|x_\gamma F_{x_\gamma}|^2 + |y_\gamma F_{y_\gamma}|^2}, \quad B := \frac{\bar{y}_\gamma \bar{F}_{y_\gamma} F_{t_\gamma}}{|x_\gamma F_{x_\gamma}|^2 + |y_\gamma F_{y_\gamma}|^2}, \quad (8.2)$$

and, by the Chain Rule,

$$\frac{\partial}{\partial x_\gamma} = \frac{\partial}{\partial x}, \quad \frac{\partial}{\partial y_\gamma} = \frac{\partial}{\partial y} + \frac{\partial \gamma}{\partial y} \frac{\partial}{\partial x}, \quad \frac{\partial}{\partial t_\gamma} = \frac{\partial}{\partial t} + \frac{\partial \gamma}{\partial t} \frac{\partial}{\partial x}. \quad (8.3)$$

The coefficients A , B are chosen so that $\langle \vec{V}, \text{Grad } F \rangle = 0$. Hence $\vec{V}$ is *tangent* to the level surfaces $F = \text{const.}$ (Here $\text{Grad } F := \bar{F}_{x_\gamma} \frac{\partial}{\partial x_\gamma} + \bar{F}_{y_\gamma} \frac{\partial}{\partial y_\gamma} + \bar{F}_{t_\gamma} \frac{\partial}{\partial t_\gamma}$.)

We have not defined $\vec{V}$ when $x_\gamma F_{x_\gamma} = y_\gamma F_{y_\gamma} = 0$. *This we shall do in §9.*

Using (8.3) we can express $\vec{V}$ as a vector field in the (x, y, t) -space:

$$\vec{\mathcal{F}}_\gamma(x, y, t) := (d\mathcal{D})^{-1}(\vec{V}), \quad (8.4)$$

which is tangent to $F = \text{const.}$ Each trajectory (integral curve) lies on a single level surface.

When $x_\gamma = 0$, the $\partial/\partial x_\gamma$ component of $\vec{V}$ vanishes, hence the flow generated by $\vec{V}$ carries the y_γ -axis to itself (but not necessarily point-wise fixed). *The flow generated by $\vec{\mathcal{F}}_\gamma$, in the (x, y, t) -space, carries the curve-germ $\pi_*(\gamma(y, 0))$ to $\pi_*(\gamma(y, t))$ at time t .*

Step Two. We are to define $\vec{\mathcal{F}}(x, y, t)$. Take $\gamma_{j,\phi} \in C(\text{val}_\phi)$ in (2.7). Take the deformation $\tau_t(\gamma_{j,\phi})$ in (2.8). Write the canonical coordinate simply as

$$\gamma_j(y, t) := \tau_t(\gamma_{j,\phi})(y), \quad 1 \leq j \leq p. \quad (8.5)$$

Take an integer N divisible by every $m_{\text{puis}}(\gamma_{j,\phi})$, for instance, $N := \prod_j m_{\text{puis}}(\gamma_{j,\phi})$.

Consider the substitution map

$$S_{XY} : (X, Y, T) \mapsto (x, y, t) := (X, Y^N, T), \quad (8.6)$$

and the coordinate transformation

$$X_j := X - \Gamma_j(Y, T), \quad Y_j := Y, \quad T_j := T,$$

where $\Gamma_j(Y, T) := \gamma_j(Y^N, T)$ is holomorphic. Like (8.1), we write

$$F^{(j)}(X_j, Y_j, T_j) := F(X_j + \Gamma_j, Y_j^N, T_j), \quad 1 \leq j \leq p.$$

Of course $X_j = 0$ is *smooth*, and is mapped by S_{XY} to $\pi_*(\gamma_{j,\phi})$, $1 \leq j \leq p$. (The latter may not be mutually distinct: if $\gamma_{1,\phi}(y)$, $\gamma_{2,\phi}(y)$ are conjugates, then $\pi_*(\gamma_{1,\phi}) = \pi_*(\gamma_{2,\phi})$.)

Therefore, for each j , $\vec{\mathcal{F}}_{\Gamma_j}(X, Y, T)$ is defined as in (8.4), being the vector field

$$\vec{V}_j(X_j, Y_j, T_j) := -A_j(X_j, Y_j, T_j)X_j \frac{\partial}{\partial X_j} - B_j(X_j, Y_j, T_j)Y_j \frac{\partial}{\partial Y_j} + \frac{\partial}{\partial T_j}$$

expressed in terms of (X, Y, T) , where, as in (8.2), with Convention 8.1,

$$A_j := \frac{\bar{X}_j \bar{F}_{X_j} F_{T_j}}{|X_j F_{X_j}|^2 + |Y_j F_{Y_j}|^2}, \quad B_j := \frac{\bar{Y}_j \bar{F}_{Y_j} F_{T_j}}{|X_j F_{X_j}|^2 + |Y_j F_{Y_j}|^2}.$$

Let us write $\hat{X}_k := X_1 \cdots X_{k-1} \cdot X_{k+1} \cdots X_p$, and, for $(X, Y) \neq (0, 0)$, define

$$\mathcal{P}_k := \frac{|\hat{X}_k|^2}{|\hat{X}_1|^2 + \cdots + |\hat{X}_p|^2}, \quad 1 \leq k \leq p. \quad (8.7)$$

We call $\{\mathcal{P}_k\}$ a *partition of unity*, for we have

$$\sum \mathcal{P}_k = 1, \quad \mathcal{P}_k = 1 \text{ when } X_k = 0, \quad \mathcal{P}_k = 0 \text{ when } X_j = 0, \quad j \neq k. \quad (8.8)$$

The $\mathcal{P}_k$'s are *real analytic* at every $(X, Y) \neq (0, 0)$ in the sense of Convention 1.1.

Now we use the $\mathcal{P}_k$'s to “patch up” the vectors $\vec{\mathcal{F}}_{\Gamma_j}$:

$$\vec{v}(X, Y, T) := \mathcal{P}_1 \vec{V}_1 + \cdots + \mathcal{P}_p \vec{V}_p \quad (\text{all expressed in } X, Y, T), \quad (8.9)$$

and, using the differential dS_{XY} of the substitution map S_{XY} , define

$$\vec{\mathcal{F}}(x, y, t) := dS_{XY}(\vec{v}). \quad (8.10)$$

We must show $\vec{\mathcal{F}}$ is well-defined, since S_{XY} is a many-to-one mapping.

Let $\theta := e^{2\pi\sqrt{-1}/N}$. A conjugation $y^{1/N} \mapsto \theta^j y^{1/N}$ permutes the γ_k 's, the X_k 's and the $\mathcal{P}_k$'s. Hence $\vec{v}(X, Y, T)$ is *invariant under these conjugations*. It follows that $\vec{\mathcal{F}}$ is well-defined.

We shall show, in §9, that $\vec{\mathcal{F}}$ can be extended continuously throughout $\mathcal{N}$, so that $\vec{\mathcal{F}}(x, y, t)$ is well-defined in $\mathcal{N}$, tangent to $F = \text{const.}$ Because of (8.8), *the flow generated by $\vec{\mathcal{F}}$ carries the curve-germ $\pi_*(\gamma_j(y, 0))$ to $\pi_*(\gamma_j(y, t))$ at time t , $1 \leq j \leq p$.*

Attention 8.2. *It is important to point out what the above does not say.*

Take $\gamma_{j,\phi}$. Of course there exists $\gamma_j(y) := \gamma_{j,\phi}(y) + \cdots$ such that $F_x(\gamma_j(y), y, 0) = 0$. Hence $F_x(x, y, 0) = 0$ on the curve-germ $\Delta := \pi_*(\gamma_j(y))$; Δ is called a “polar” of $F(x, y, 0)$.

Note that $\mathcal{C}_{ord}(\Delta, \pi_*(\gamma_{j,\phi})) > O_y(\gamma_{j,\phi})$, and, in general, $\Delta \neq \pi_*(\gamma_{j,\phi})$. Following the flow, Δ reaches Δ_t at time t . *The above does not say Δ_t is necessarily a polar of $F(x, y, t)$.*

For example, the Pham family $P(x, y, t)$ in (0.1) has only *one* polar when $t = 0$, but *two* polars when $t \neq 0$. We have no idea whether the polar at $t = 0$ will flow to one of the two polars, or more likely to neither. In the *blurred space* $\mathbb{C}_*/P_t$, however, there is a *unique critical point* for each $t \in I_{\mathbb{C}}$; they constitute a single orbit of $\{\eta_t\}$.

A critical point is an equivalence class in $\mathbb{C}_*$ containing at least one polar. *We see the critical point at all time t , but cannot keep track of the polars.* (This is like the Arakawa in Japan, a river which flows by Saitama University. We see the river bed, but cannot predict the position of the flow, whence, literally, the name “Arakawa”–Wild River.)

9. PROOF OF THE EQUI-SINGULAR DEFORMATION THEOREM

Recall that F is mini-regular in x . In a sector $|y| \leq \epsilon|x|$, the behavior of F is dominated by x^m , $m := O(F)$. Hence there is nothing to worry about in this sector.

We shall henceforth restrict our attention to a sector $|x| < K|y|$, K sufficiently large.

Notation 9.1. In this section we write $g \lesssim h$ if $g \leq Ch$, $C > 0$ a constant; $g \approx h$ means $g \lesssim h \lesssim g$; and $g \ll h$ means $g/h \rightarrow 0$.

Next we show how $\vec{v}$ in (8.9) and $\vec{\mathcal{F}}$ in (8.10) generate homeomorphisms.

The following is a parameterized version of the Proposition in [18], p.347.

Lemma 9.2. For $F^{(j)}(X_j, Y_j, T_j) := F(X_j + \Gamma_j(Y_j, T_j), Y_j^N, T_j)$, we have

$$|F_{T_j}| \lesssim |X_j F_{X_j}| + |Y_j F_{Y_j}|, \quad (X_j, Y_j, T_j) \in S_{XY}^{-1}(\mathcal{N}), \quad (9.1)$$

where $1 \leq j \leq p$, $\mathcal{N}$ as in §8. (We use Convention (8.1): $F_{X_j} := F_{X_j}^{(j)}$, etc..)

We can extend A_j , B_j real analytically to $S_{XY}^{-1}(\mathcal{N}) - \{0\} \times I_{\mathbb{C}}$, where they are bounded.

Define $\vec{\mathcal{F}}(0, 0, t) := \frac{\partial}{\partial t}$. Then $\vec{\mathcal{F}}$ is continuous on $\mathcal{N}$,

$$\|\vec{\mathcal{F}}(x, y, t) - \partial/\partial t\| \lesssim |x| + |y|, \quad (x, y, t) \in \mathcal{N}, \quad (9.2)$$

and $\vec{\mathcal{F}}$ is real analytic in $\mathcal{N} - \{0\} \times I_{\mathbb{C}}$.

It follows that a trajectory of $\vec{\mathcal{F}}$, with initial point outside $I_{\mathbb{C}}$, will never reach $I_{\mathbb{C}}$.

The flow generated by $\vec{\mathcal{F}}$ carries $\pi_*(\gamma_{j,\phi})$ to $\pi_*(\tau_t(\gamma_{j,\phi}))$, $1 \leq j \leq p$.

Proof. We use the Curve Selection Lemma to prove (9.1). Let $\rho(s)$ be a given analytic arc. It suffices to show that (9.1) holds along ρ . We can assume $\rho(0) = 0$.

Take $\epsilon' > 0$, sufficiently small. For each pair Γ_k, Γ_s (in §8), let $d_{ks} := O_Y(\Gamma_k - \Gamma_s)$.

Define a horn neighborhood of Γ_k of order d_{ks} by

$$H_{d_{ks}}(\Gamma_k) := \{(X, Y, T) \mid |X - \Gamma_k(Y, T)| < \epsilon'|Y|^{d_{ks}}\}. \quad (9.3)$$

This is a sub-analytic set, hence either $Im(\rho) - \{0\} \subset H_{d_{ks}}(\Gamma_k)$ or $Im(\rho) \cap H_{d_{ks}}(\Gamma_k) = \emptyset$.

Let us first consider the case where $Im(\rho) - \{0\}$ is contained in at least one of the horn neighborhoods. In this case, by permuting the indices, if necessary, we can assume $H_{d_{12}}(\Gamma_1)$ is the *smallest* horn neighborhood containing $Im(\rho) - \{0\}$.

We then work in the coordinate system (X_1, Y_1, T_1) , writing $\rho(s) = (X_1(s), Y_1(s), T_1(s))$.

Let $\rho_\pi(x) := (X_1(s), Y_1(s))$. We show (9.1) holds on the surface $Im(\rho_\pi) \times I_{\mathbb{C}}$.

Let us write

$$F^{(1)}(X_1, Y_1, T_1) := F_0^{(1)}(X_1, Y_1) + P(X_1, Y_1, T_1), \quad P(X_1, Y_1, 0) \equiv 0. \quad (9.4)$$

By the Fundamental Lemma, $\mathcal{NP}(F^{(1)}, 0) = \mathcal{NP}(F_0^{(1)}, 0)$, all dots of $P(X_1, Y_1, T_1)$ lie on or above this polygon. (Newton Polygon at 0 is Newton Polygon in the usual sense.)

Convention 9.3. Consider the vertex $V_l = (m_l, q_l)$. Suppose $m_l = 0$ and $h > \tan \theta_{top}$. In this section, we call the pair $E(h) := (V_l, h)$ also a *vertex edge*, with co-slope h .

First, suppose $Im(\rho)$ is not contained in $X_1 = 0$. We write the coordinate of $\rho_{\pi, \mathbb{C}}$ as

$$\rho_{\pi}(Y_1) := rY_1^h + \cdots, \quad r \neq 0, \quad d_{12} \leq h < \infty. \quad (9.5)$$

And let $E := E(h)$ denote the *unique* (possibly vertex) edge with co-slope h .

We first prove (9.1) for $j = 1$.

Assume $E := E(h)$ is a proper edge (*i.e.*, not a vertex edge).

In the first place we must have $P'_E(r) \neq 0$. For if $P'_E(r) = 0$, then, by Theorem 5.3, there would exist Γ_j , $j \geq 2$, of the form $\Gamma_j = rY_1^h + \cdots$. Then $H_{d_{12}}(\Gamma_1)$ would not be the smallest horn neighbourhood containing $Im(\rho)$, a contradiction.

Now we collect the monomial terms of $F^{(1)}$ along E :

$$W(X_1, Y_1, T_1) := \sum a_{pq}(t) X_1^p Y_1^q, \quad (p, q) \in E, \quad (9.6)$$

which is a weighted form, $W(z, 1, t) = P_E(z)$.

Take $u \neq 0$. Let t be fixed. By Euler's Theorem, $X_1 - uY_1^h$ is a common factor of $X_1 W_{X_1}$ and $Y_1 W_{Y_1}$ iff $(X_1 - uY_1^h)^2$ divides $W(X_1, Y_1, T_1)$, *i.e.*, $P_E(u) = P'_E(u) = 0$.

Hence, if $(P_E(r), P'_E(r)) \neq (0, 0)$, then, along ρ_{π} ,

$$O_{Y_1}(|X_1 W_{X_1}| + |Y_1 W_{Y_1}|) = O_{Y_1}(|X_1 F_{X_1}| + |Y_1 F_{Y_1}|) = q_E + m_E h,$$

where (m_E, q_E) is any dot on E . It follows that

$$L_{|X_1 F_{X_1}| + |Y_1 F_{Y_1}|}(\rho_{\pi}) = q_E + m_E h. \quad (9.7)$$

All dots of F_{T_1} lie on or above the line $\mathcal{L}(E)$, $O_{Y_1}(F_{T_1}) \geq q_E + m_E h$, proving (9.1).

If $E = (V_k, h)$ is a vertex edge, then $|Y_1 F_{Y_1}| \approx |r|^{m_k} |Y_1|^{q_k + m_k h} \gtrsim |F_{T_1}|$. Again (9.1) holds.

Next we prove (9.1) for the case $j \geq 2$. The coordinate systems are related by

$$X_j = X_1 - \delta_j(Y_1, T_1), \quad Y_j = Y_1, \quad T_j = T_1, \quad (9.8)$$

where $\delta_j := \Gamma_j - \Gamma_1$ ($\delta_1 \equiv 0$). Let us write

$$\delta_j(Y_1, T_1) := c_j(T_1) Y_1^{O(\delta_j)} + \cdots, \quad c_j(0) \neq 0, \quad 2 \leq j \leq p. \quad (9.9)$$

Note that

$$|Y_1 \frac{\partial \delta_j}{\partial Y_1}| \approx |\delta_j|, \quad |\frac{\partial \delta_j}{\partial T_1}| \lesssim |\delta_j|,$$

and, by the Chain Rule,

$$X_j F_{X_j} = (X_1 - \delta_j) F_{X_1}, \quad Y_j F_{Y_j} = Y_1 F_{Y_1} + Y_1 \frac{\partial \delta_j}{\partial Y_1} F_{X_1}, \quad F_{T_j} = F_{T_1} + \frac{\partial \delta_j}{\partial T_1} F_{X_1}. \quad (9.10)$$

Now, if $h \geq O(\delta_j)$, then, along ρ_{π} , $|X_1 - \delta_j| \approx |\delta_j|$. Hence (9.1) follows from

$$|X_j F_{X_j}| + |Y_j F_{Y_j}| \approx |\delta_j F_{X_1}| + |Y_1 F_{Y_1}|, \quad |F_{T_j}| \lesssim |F_{T_1}| + |\delta_j F_{X_1}|,$$

and (9.1) when $j = 1$.

Suppose $h < O(\delta_j)$. Then $|X_1 - \delta_j| \approx |X_1|$ along ρ_{π} . Again we have (9.1).

Suppose $Im(\rho)$ is contained in $X_1 = 0$. In this case F is divisible by $X_1^{m_l}$, F_{T_1} is divisible by X_1^k , $k \geq m_l$. Again, we have (9.1).

It remains to consider the case where no horn neighborhood contains $Im(\rho) - \{0\}$. We again write ρ_π as (9.5) where now $h < d_{jk} \forall j, k$. The same argument proves (9.1).

Next we show how A_j, B_j can be extended across $S_{XY}^{-1}(\mathcal{N}) - \{0\} \times I_{\mathbb{C}}$.

Lemma 9.4. *The zero set of the denominator of A_j, B_j is*

$$\{(X, Y, T) | X_j F_{X_j} = Y_j F_{Y_j} = 0\} = \begin{cases} \bigcup \{X_k = 0 \mid X_k^2 \text{ divides } F\}, \\ \{0\} \quad \text{if no such } X_k \text{ exists.} \end{cases}$$

Proof. Recall that we work in the sector $|x| < K|y|$, hence $Y \neq 0$.

Suppose $X_j = F_{Y_j} = 0$. In $\mathcal{NP}(F^{(j)}, \Gamma_j)$, which is independent of T_j , we must have $m_l \geq 2$. Hence F is divisible by X_j^2 .

Let $\rho(s)$ be an analytic arc along which $F_{X_j} = F_{Y_j} = 0$. Then $F_X = F_Y = 0$ along ρ . As before, we choose X_1 , define ρ_π , $F_0(X_1, Y_1)$, etc., as in (9.4), and then

$$\frac{\partial F_0^{(1)}}{\partial X_1} = \frac{\partial F_0^{(1)}}{\partial Y_1} = 0 \quad \text{along } \rho_\pi.$$

Hence $Im(\rho_{\pi, \mathbb{C}})$ must be $X_1 = 0$ ($T_1 = 0$), X_1^2 divides F_0 , hence also F . $\square$

We can now complete the proof of Lemma 9.2.

The real meromorphic functions A_j, B_j are *bounded*, by (9.1). Hence, if n is the largest integer such that X_k^n divides both $X_j F_{X_j}$ and $Y_j F_{Y_j}$, then X_k^n must also divide F_{T_j} .

It follows that A_j, B_j are *defined and real analytic* on $S_{XY}^{-1}(\mathcal{N}) - \{0\} \times I_{\mathbb{C}}$.

Hence $\vec{\mathcal{F}}(x, y, t)$, with $\vec{\mathcal{F}}(0, 0, t) := \frac{\partial}{\partial t}$, is *continuous* in $\mathcal{N}$, *satisfying* (9.2), and is *real analytic* in $\mathcal{N} - \{0\} \times I_{\mathbb{C}}$.

Finally, $\mathcal{P}_k X_k = 0$ along every $X_j = 0$. Hence $\vec{\mathcal{F}}$ carries $\pi_*(\gamma_{j, \phi})$ to $\pi_*(\tau_t(\gamma_{i, \phi}))$. $\square$

Now, using a well-known argument ([12]), (9.2) implies that $\vec{\mathcal{F}}$ generates a homeomorphism H having properties (3.1), (1) and (2) in the Equi-singular Deformation Theorem.

Next we prove (3). Let $\rho(s) = (x(s), y(s))$ be a given analytic arc in the (x, y) -plane.

Consider $\rho^{(N)}(s) := (x(s^N), y(s^N))$. This arc can be lifted by S_{XY} to an analytic arc

$$\rho_{XY}(s) := (X(s), Y(s)), \quad S_{XY} \circ \rho_{XY} = \rho^{(N)},$$

where $Y(s)$ is an *integral* power series in s obtained by solving $Y^N = y(s^N)$. (Of course, there are N liftings of $\rho^{(N)}$ to the (X, Y) -space; in general, however, we cannot lift ρ .)

Let us write $\Gamma_j := \Gamma_j(Y, 0)$. By permuting the indices, if necessary, we can assume

$$\begin{aligned} h &:= \mathcal{C}_{ord}(Im(\rho_{XY, \mathbb{C}}), Im(\Gamma_1)) = \dots = \mathcal{C}_{ord}(Im(\rho_{XY, \mathbb{C}}), Im(\Gamma_r)) \\ &> \mathcal{C}_{ord}(Im(\rho_{XY, \mathbb{C}}), Im(\Gamma_{r+i})), \quad 0 < i \leq p - r, \end{aligned} \tag{9.11}$$

where $1 \leq r \leq p$, $h/N = \mathcal{C}_{ord}(Im(\rho_{\mathbb{C}}), \pi_*(\gamma_{j, \phi}))$, $1 \leq j \leq r$.

There are now three cases to consider: (a) $h/N \leq h(\gamma_{1,\phi})$, $h < \infty$, (b) $h(\gamma_{1,\phi}) < h/N < \infty$, and (c) $h = \infty$, that is, $Im(\rho_c) = \pi_*(\gamma_{1,\phi})$.

Case (c) is easy: $Im(\rho_{XY,\mathbb{C}})$ is the Y_1 -axis, along which $\vec{v} = \vec{V}_1$ is analytic, whence (3).

Consider case (a). Let $E := E(h)$ be the unique edge in $\mathcal{NP}_{ext}(F_0^{(1)}, 0)$ of co-slope h . (Of course, E can be a vertex edge (V_e, h) . In this case $m_e > 0$, $P_E(z)$ is a monomial.)

As before, we work in the coordinate system (X_1, Y_1, T_1) .

Take a coordinate of $Im(\rho_{XY,\mathbb{C}})$,

$$\rho_{XY}(Y_1) := u_\rho Y_1^h + \cdots, \quad u_\rho \neq 0.$$

A coordinate of $Im(\rho_c)$ is

$$\rho(y) = \gamma_{1,\phi}(y) + [u_\rho y^{h/N} + \cdots].$$

As before, $P'_E(u_\rho) \neq 0$. (Same argument: otherwise, (9.11) would fail.)

Now, let us first assume that h is an integer, so that the substitution map

$$S_{uv} : (u, v, t) \mapsto (X_1, Y_1, T_1) := (uv^h, v, t),$$

is holomorphic, where

$$X_1 \frac{\partial}{\partial X_1} = u \frac{\partial}{\partial u}, \quad Y_1 \frac{\partial}{\partial Y_1} = -h u \frac{\partial}{\partial u} + v \frac{\partial}{\partial v}, \quad \frac{\partial}{\partial T_1} = \frac{\partial}{\partial t}. \quad (9.12)$$

We then define

$$\vec{U}(u, v, t) := (dS_{uv})^{-1}(\vec{v}).$$

Question 9.5. *At which (u, v, t) is $\vec{U}$ well-defined and real analytic?*

To answer this, we need a careful analysis of the denominator of A_j, B_j :

$$D_j := |X_j F_{X_j}|^2 + |Y_j F_{Y_j}|^2, \quad 1 \leq j \leq p,$$

and also that of $\mathcal{P}_k$, when the substitution S_{uv} is made.

Lemma 9.6. *Let $L(E) := q_E + m_E h$, $(m_E, q_E) \in E$. Then*

$$D_j(uv^h, v, t) = C_j(u, t) v^{2L(E)} + \cdots, \quad 1 \leq j \leq r, \quad (9.13)$$

where $C_j(u, t)$ is a polynomial in $u, \bar{u}$, coefficients in t .

For $u \neq 0$, $C_j(u, t) = 0$ iff $P_E(u) = P'_E(u) = 0$. For any u , $P'_E(u) \neq 0$ implies $C_j(u, t) \neq 0$.

(The coefficients of $P_E(u)$ are functions of t . To say $P_E(u) = P'_E(u) = 0$ means that when t is fixed, u is a multiple root of $P_E(z)$.)

Proof. Consider the weighted form $W(X_1, Y_1, T_1)$ in (9.6). Let

$$w_1(z, t) := W_{X_1}(z, 1, t), \quad w_2(z, t) := W_{Y_1}(z, 1, t).$$

Take j , $1 \leq j \leq r$. By (9.11), $O(\delta_j) \geq h$, δ_j being defined in (9.8). Let us write

$$\hat{c}_j(t) := \begin{cases} c_j(t) & \text{if } O(\delta_j) = h, \\ 0 & \text{if } O(\delta_j) > h, \end{cases}$$

where $c_j(t)$ is defined in (9.9). We claim that (9.13) holds if we take

$$C_j(u, t) := |(u - \hat{c}_j(t))w_1(u, t)|^2 + |w_2(u, t) + h\hat{c}_j(t)w_1(u, t)|^2,$$

which is of course a polynomial in $u, \bar{u}$. Indeed, we have, as before,

$$D_j = |(X_1 - \delta_j)F_{X_1}|^2 + |Y_1 F_{Y_1} + Y_1 \frac{\partial \delta_j}{\partial Y_1} F_{X_1}|^2, \quad (9.14)$$

whence the leading term of $D_j(uv^h, v, t)$ is the above $C_j(u, t)$.

The $\hat{c}_i(t)$'s are roots of $w_1(z, t) = P'_E(z)$. Hence $C_j(u, t) = 0$ iff $w_1(u, t) = w_2(u, t) = 0$.

For $u \neq 0$, $w_1 = w_2 = 0$ iff $(X_1 - uY_1^h)^2$ divides $W(X_1, Y_1, T_1)$, i.e., $P_E(u) = P'_E(u) = 0$. But we already know $P'_E(0) = 0$. Hence $P'_E(u) \neq 0$ implies $u \neq 0$, $C_j(u, t) \neq 0$. $\square$

Lemma 9.7. *For $r < j \leq p$, we have*

$$D_j(uv^h, v, t) = (1 + h^2)|c_j(t)w_1(u, t)|^2 v^{2[L(E) - h + O_y(\delta_j)]} + \dots \quad (9.15)$$

Hence if $P'_E(u) \neq 0$, then $O_v(D_j(uv^h, v, t)) = 2[L(E) - h + O_y(\delta_j)]$.

If $P_E(u) \neq 0$, then $O_v(D_j(uv^h, v, t)) \leq 2L(E)$.

Proof. If $j > r$, then $h > O(\delta_j)$ and hence

$$uv^h - \delta_j = -c_j(t)v^{O(\delta_j)} + \dots, \quad c_j(0) \neq 0.$$

Then, as can be observed from (9.14), the leading term of D_j is that of (9.15).

Next note that

$$D_j \approx |\delta_j F_{X_1}|^2 + |v F_{Y_1} + v \frac{\partial \delta_j}{\partial Y_1} F_{X_1}|^2 \approx |\delta_j F_{X_1}|^2 + |v F_{Y_1}|^2 \geq |v F_{Y_1}|^2,$$

and that $O_v(v F_{Y_1}(uv^r, v, t)) = L(E)$ if $P_E(u) \neq 0$. This completes the proof. $\square$

The Newton dots of F_{T_j} lie on or above the line $\mathcal{L}(E)$, hence the above lemmas imply that $A_j \circ S_{uv}$, $B_j \circ S_{uv}$ are real analytic at (u, v, t) if $P'_E(u) \neq 0$, $1 \leq j \leq p$.

Lemma 9.8. *The denominator $D_{\mathcal{P}} := |\hat{X}_1|^2 + \dots + |\hat{X}_p|^2$ of $\mathcal{P}_k$ has the form*

$$D_{\mathcal{P}}(uv^h, v, t) = c(u, t)v^{2e} + \dots, \quad e := (r - 1)h + \sum_{j>r} O(\delta_j), \quad (9.16)$$

where $c(u, t)$ is a polynomial in $u, \bar{u}$.

If $(P_E(u), P'_E(u)) \neq (0, 0)$ then $c(u, t) \neq 0$.

Proof. For $1 \leq j \leq r$, let us write

$$\mu_j(u, t) := u - \hat{c}_j(t), \quad \hat{\mu}_j(u, t) := \mu_1 \cdots \mu_{j-1} \cdot \mu_{j+1} \cdots \mu_r.$$

We can compute the $\hat{X}_k$'s, $1 \leq k \leq p$, using the formula

$$uv^h - \delta_k = \begin{cases} (u - \hat{c}_k(t))v^h + \dots, & \text{if } 1 \leq k \leq r, \\ -c_k(t)v^{O(\delta_k)} + \dots, & \text{if } r < k \leq p, \end{cases}$$

where $c_k(0) \neq 0$ for $r < k \leq p$. For example,

$$\hat{X}_1 = [(-1)^{p-r} c_{r+1}(t) \cdots c_p(t)](u - \hat{c}_2(t)) \cdots (u - \hat{c}_r(t))v^e + \dots,$$

where e is defined in (9.16). The equations for the other $\hat{X}_k$'s are similar.

Then (9.16) holds when we take

$$c(u, t) := |c_{r+1}(t) \cdots c_p(t)|^2 \cdot \sum_{j=1}^r |\hat{\mu}_j(u, t)|^2.$$

If $c(u, t) = 0$, then all $\hat{\mu}_j = 0$, and there exist i, j , $1 \leq i < j \leq r$, $u = \hat{c}_i(t) = \hat{c}_j(t)$. This implies, in particular, that $P'_E(u) = 0$.

We claim this also implies $P_E(u) = 0$. Indeed, $\hat{c}_i(t) = \hat{c}_j(t)$ implies $O(\gamma_{i,\phi} - \gamma_{j,\phi}) > h/N$. If $P_E(u) \neq 0$, then $h/N = h(\gamma_{i,\phi}) = h(\gamma_{j,\phi})$, hence $\gamma_{i,\phi} = \gamma_{j,\phi}$, a contradiction. $\square$

The $\mathcal{P}_k$'s are bounded. Hence $\mathcal{P}_k \circ S_{uv}$ are real analytic at (u, t) if $(P_E(u), P'_E(u)) \neq (0, 0)$.

Let $\mathcal{N}'$ be an open neighborhood of the u -axis such that $(u, v) \in \mathcal{N}'$ implies $(uv^r, v) \in \mathcal{N}$.

Answer to Question 9.5: Take $(u_0, v_0, t_0) \in \mathcal{N}' \times I_{\mathbb{C}}$. If (u_0, t_0) is not a root of $P'_E(u)$, then $\vec{U}(u, v, t)$ is defined and real analytic in a neighborhood of (u_0, v_0, t_0) .

Lemma 9.9. Given j , $1 \leq j \leq r$. For (u, t) near $(\hat{c}_j(t), t)$,

$$|\vec{U}(u, v, t) - \frac{\partial}{\partial t}| \lesssim |u - \hat{c}_j(t)| + |v|. \quad (9.17)$$

Moreover, $\vec{U}(u, 0, t)$ is tangent to the u -coordinate space, and

$$|\vec{U}(u, 0, t) - \frac{\partial}{\partial t}| \lesssim |u - \hat{c}_j(t)|. \quad (9.18)$$

Proof. Let us first assume $j = 1$, where $\hat{c}_1(t) \equiv 0$. The notations are simpler.

First, by (9.12),

$$(dS_{uv})^{-1}(\mathcal{P}_1 \vec{V}_1) = -\mathcal{P}_1 A_1 u \frac{\partial}{\partial u} - \mathcal{P}_1 B_1 (-hu \frac{\partial}{\partial u} + v \frac{\partial}{\partial v}) + \mathcal{P}_1 \frac{\partial}{\partial t},$$

where $\mathcal{P}_1 A_1, \mathcal{P}_1 B_1$ are bounded.

For k , $2 \leq k \leq p$, the identity

$$\mathcal{P}_k X_k \frac{\partial}{\partial X_k} = \{\mathcal{P}_k^{1/2} \mathcal{P}_1^{1/2} X_k |X_k|^{-1}\} |X_1| \frac{\partial}{\partial X_1},$$

and the Chain Rule (9.12) give

$$(dS_{uv})^{-1}(\mathcal{P}_k \vec{V}_k) = \tilde{A} u \frac{\partial}{\partial u} + \tilde{B} v \frac{\partial}{\partial v} + \frac{\partial}{\partial t},$$

where $\tilde{A}, \tilde{B}$ are bounded.

Hence (9.17) is true for $j = 1$.

Now assume $1 < j \leq r$. This case is actually the same as the case $j = 1$. For in (9.11), the roles of Γ_1 and Γ_j are interchangeable. When Γ_j replaces Γ_1 , the leading coefficient $c_k(t)$ of δ_k in (9.9) is replaced by $c_k(t) - c_j(t)$. The same argument completes the proof of (9.17).

The $\frac{\partial}{\partial v}$ component vanishes when $v = 0$. Hence $\vec{U}(u, 0, t)$ is tangent to the u -space. $\square$

We are now ready to complete the proof of (3) in the case (a).

Recall that $Im(\Gamma_1)$ is the vertical axis $X_1 = 0$, and $\rho_{XY,C}$ is defined by $X_1 = u_\rho Y_1^h + \dots$, which lifts to $\rho_{uv} : u = u_\rho + \dots$.

We know $P'_E(u_\rho) \neq 0$. Hence the arc ρ_{uv} lies in the domain where $\vec{U}$ is analytic. The flow carries ρ_{uv} to an analytic arc, at least for a sufficiently short time. Hence the flow of $\vec{v}$ carries ρ_{XY} to an analytic arc, at least for a short time. (Keep $|v|$ small, arcs short.)

This is actually so for *all* $t \in I_C$, not just for a short time. Indeed, using Lemma 9.9, a well-known argument shows that a point on ρ_{uv} , following the flow, will *never* reach a point where $\vec{U}$ is not analytic, whence the trajectory is defined for *all* t .

The initial point $(u_\rho, 0, 0)$ of ρ_{uv} *remains* in the u -coordinate space for all time, because $\vec{U}$ is tangent to the u -space. By (9.18), it will never reach $\hat{c}_j(t)$.

Hence, downstairs, the geo-arc $Im(\rho)$ is carried by the flow of $\vec{F}$ real analytically, sweeping out a “geo-arc wing”. This completes the proof of (3) in the case (a) when $h \in \mathbb{Z}^+$.

Suppose $h := N_1/M_1 \notin \mathbb{Z}^+$. We can make a further substitution $Y_1 \mapsto Y_1^{M_1}$. The vector field lifts, retaining the same form, h is magnified to N_1 . The same argument applies.

Finally, consider case (b). This can be treated as a special case of (a), as follows.

In $\mathcal{NP}(\phi, \gamma_{1,\phi})$, E_{top} has left vertex $V_l = (0, q_l)$, since $h(\gamma_{1,\phi}) < \infty$. We may call $E := (V_l, h)$ an “artificial vertex edge”, of co-slope h , with Lojasiewicz exponent $L(E) := q_l$.

Let V_l represent the term cy^{q_l} , $c \neq 0$. The *associated polynomial* is $P_E(z) := c$.

The above argument for the case (a) can then be repeated. It is actually easier. The lemmas remain true for the artificial vertex edge E , where now $P_E(u) = c \neq 0$.

This completes the proof of (3).

We then use the same idea to prove (4).

Take $\alpha_{*/f} \in \mathbb{C}_{*/f}$ and a canonical coordinate $\alpha_\phi(y)$. We can assume

$$h(\alpha_\phi) = O_y(\alpha_\phi - \zeta_1) \geq O_y(\alpha_\phi - \zeta_i), \quad O_y(\alpha_\phi - \gamma_{1,\phi}) \geq O_y(\alpha_\phi - \gamma_{j,\phi}), \quad (9.19)$$

where $\zeta_i \in Z(\phi)$, $\gamma_{j,\phi} \in C(val_\phi)$.

We now repeat the argument in (3) to show $\eta_t(\alpha_{*/f})$ is well-defined. Let us write

$$\alpha_\phi(y) := \zeta_1(y) + [u_0 y^{h(\alpha_\phi)} + \dots], \quad u_0 \neq 0, \quad (9.20)$$

where $\alpha_{*/f}$ is *completely* determined by u_0 . (Terms in “ $+\dots$ ” play no role.)

Here we have assumed that $h(\alpha_\phi) < \infty$. (If $\alpha_\phi = \zeta_1$, we can apply Corollary 5.6.)

Take an analytic arc ρ such that $\pi_*(\rho_c) \in \alpha_{*/f}$. As in (3), we lift ρ to ρ_{XY} , and analyze how ρ_{XY} is being carried by $\vec{v}(X, Y, T)$ in (8.9).

This time we use the coordinate system:

$$X_\zeta := X - \zeta_1(Y^N, T), \quad Y_\zeta := Y, \quad T_\zeta := T, \quad (y := Y^N),$$

where N is divisible by $m_{puls}(\alpha_\phi)$ and $m_{puls}(\zeta_i)$.

The substitution $(X_\zeta, Y_\zeta) := (uv^h, v)$, $h := Nh(\alpha_\phi)$, lifts ρ_{XY} to an analytic arc ρ_{uv} in the (u, v) -space whose initial point is $(u_0, 0)$.

As in (3), $\vec{v}$ is lifted to $\vec{U}$; ρ_{uv} lies in the domain where $\vec{U}$ is real analytic.

Hence, following the flow of $\vec{U}$ for time t , ρ_{uv} reaches an analytic arc, denoted by $\rho_{uv,t}$, with initial point $(u_t, 0, t)$, where u_t is real analytic in t .

An important observation is that $(u_0, 0)$, and hence also $(u_t, 0, t)$, depend *only* on $\alpha_{*/f}$, *not* on the choice of ρ . It follows that $\eta_t(\alpha_{*/f})$ is well-defined.

(Note. Even if ρ, μ have the *same* complexification, $Im(\rho_c) = Im(\mu_c)$, the above argument *does not* prove that the geo-arcs $Im(\eta_t \circ \rho)$, $Im(\eta_t \circ \mu)$ lie on a *same* curve-germ.

It *merely* shows that the curve-germs containing $Im(\eta_t \circ \rho)$ and $Im(\eta_t \circ \mu)$ are equivalent under $\sim_{F_t}$. This is the ‘‘Arakawa phenomenon’’: *only the blurred point is well-defined.*)

To complete the proof of (4), we need to know how to obtain $\mathcal{NP}(f, \alpha_\phi)$ from $\mathcal{NP}(f, \zeta_1)$.

If $h(\alpha_\phi) = \infty$, the two polygons are identical.

Assume $h(\alpha_\phi) < \infty$ ($u_0 \neq 0$) in (9.20). Let $\{(m_1, q_1), \dots, (m_l, q_l)\}$ denote the vertices of $\mathcal{NP}(f, \zeta_1)$, where, of course, $m_l \geq 1$.

Take k ($k \leq l$) such that $\tan \theta_{k-1} \leq h(\alpha_\phi) < \tan \theta_k$.

Let $(0, q')$ be the left vertex of the edge $E(h(\alpha_\phi))$. (If $\tan \theta_{k-1} < h(\alpha_\phi) < \tan \theta_k$, then $E(h(\alpha_\phi))$ is a vertex edge.) The vertices of $\mathcal{NP}(f, \alpha_\phi)$ are

$$\begin{cases} (m_1, q_1), \dots, (m_{k-1}, q_{k-1}), (0, q'), & \text{if } \tan \theta_{k-1} = h(\alpha_\phi), \\ (m_1, q_1), \dots, (m_l, q_l), (0, q'), & \text{if } \tan \theta_{k-1} < h(\alpha_\phi). \end{cases}$$

We can also obtain $\mathcal{NP}(F_t, \eta_t(\alpha_{*/f}))$ from $\mathcal{NP}(F_t, \zeta_{1,t})$ in the same way, since $u_t \neq 0$.

By Corollary 5.6, we know $\mathcal{NP}(F_t, \zeta_{1,t}) = \mathcal{NP}(f, \zeta_1)$. Hence $\mathcal{NP}(F_t, \eta_t(\alpha_{*/f})) = \mathcal{NP}(f, \alpha_\phi)$. It follows that $h(\eta_t(\alpha_{*/f}))$, $\chi_{\text{puis}}(\eta_t(\alpha_{*/f}))$ are constants.

It remains to consider the contact order. From what we have proved,

$$\mathcal{C}_{ord}(\alpha_{*/f}, \pi_*(\zeta_i)) = \mathcal{C}_{ord}(\eta_t(\alpha_{*/f}), \pi_*(\zeta_{i,t})), \quad \zeta_i \in Z(\phi).$$

And, by Corollary 9.20,

$$\mathcal{C}_{ord}(\pi_*(\zeta_{i,t}), \pi_*(\zeta_{j,t})) = \mathcal{C}_{ord}(\pi_*(\zeta_i), \pi_*(\zeta_j)), \quad \zeta_i, \zeta_j \in Z(\phi).$$

The same holds for $\beta_{*/f}$. It follows that $\mathcal{C}_{ord}(\eta_t(\alpha_{*/f}), \eta_t(\beta_{*/f}))$ is independent of t . This completes the proof of (4).

The vector field $\vec{\mathcal{F}}$ is defined in such a way that (5) is true.

Finally, suppose Φ is Morse stable. Then (3.3) is a consequence of (2.9). This completes the proof of the Equi-singular Deformation Theorem.

To prove the Truncation Theorem, note that $\mathcal{NP}(f, \alpha_\phi)$ and $\mathcal{NP}(u \cdot f, \alpha_\phi)$ have the *same* Newton dots, where $u(0, 0) \neq 0$. (The dots in the interior of the polygons may be different.)

The monomial terms of $\mathcal{NP}(u \cdot f, \alpha_\phi)$ are those of $\mathcal{NP}(f, \alpha_\phi)$ multiplied by the *same* constant $u(0, 0)$. Hence the critical points $C(val_\phi)$ is *unchanged* when f is multiplied by a unit, F_{root} is *obviously* Morse stable. Now apply the Equi-singular Deformation Theorem.

10. APPENDIX 1. THE STABILITY THEOREM

The Classical Morse Stability Theorem. *A Morse stable family $\{p_t(x)\}$, as in Definition 5.4, $\mathbb{K} = \mathbb{R}, \mathbb{C}$, or $\mathbb{F}$, admits a real-analytic trivialization. That is to say, there exist t -level preserving, real-analytic, homeomorphisms (deformations) $\mathcal{D}$ and d , such that*

$$\begin{array}{ccc} \mathbb{K} \times I_{\mathbb{K}} & \xrightarrow{g} & \mathbb{K} \times I_{\mathbb{K}} \\ \downarrow \mathcal{D} & & \downarrow d \\ \mathbb{K} \times I_{\mathbb{K}} & \xrightarrow{G} & \mathbb{K} \times I_{\mathbb{K}} \end{array}$$

is commutative, where $d(0, t) = (0, t)$, $g(x, t) := (p_0(x), t)$, $G(x, t) := (p_t(x), t)$.

(In the case $\mathbb{K} = \mathbb{C}$, moduli can appear, we cannot ask $\mathcal{D}$, d to be holomorphic.)

Proof. The critical value set of G is, by definition,

$$V_{crit}(G) := \{(v_t, t) \in \mathbb{K} \times I_{\mathbb{K}} \mid \exists c_t \in \mathbb{K}, p'_t(c_t) = 0, v_t = p_t(c_t)\},$$

and $\pi : V_{crit}(G) \rightarrow I_{\mathbb{K}}$ is a fibration, $\{0\} \times I_{\mathbb{K}}$ is either disjoint from, or contained in, $V_{crit}(G)$.

Let us first consider the case $\mathbb{K} = \mathbb{R}$, which exposes the main ideas. Take a vector field

$$\vec{v}(x, t) := a(x, t) \frac{\partial}{\partial x} + \frac{\partial}{\partial t}, \quad a(x, t) \text{ analytic,}$$

which is defined on, and tangent to, $V_{crit}(G) \cup (\{0\} \times I_{\mathbb{R}})$.

Then, using Cartan's Theorem B, or the Lagrange Interpolation Formula, we can extend $a(x, t)$ to a real analytic function defined for $(x, t) \in \mathbb{R} \times I_{\mathbb{R}}$.

In other words, $\vec{v}(x, t)$ is now defined and real analytic on $\mathbb{R} \times I_{\mathbb{R}}$, tangent to $V_{crit}(G)$ and $\{0\} \times I_{\mathbb{R}}$. Integrating this vector field gives an analytic deformation

$$d : \mathbb{R} \times I_{\mathbb{R}} \rightarrow \mathbb{R} \times I_{\mathbb{R}}, \quad (x, t) \mapsto (x_t, t),$$

such that $d(V_{crit}(g)) = V_{crit}(G)$, and $\{0\} \times I_{\mathbb{R}}$ is fixed.

Using d as an identification, we assume $V_{crit}(g) = V_{crit}(G)$ ("straightening up" $V_{crit}(G)$).

Take $c \in C(p_0)$, $p_0(c) := v$. Then c admits a *unique continuous* deformation c_t in $C(p_t)$, $p_t(c_t) = v$, $c_0 = c$. As $m_{crit}(c_t)$ is constant, c_t is necessarily *real analytic*. Hence

$$p_t(x) - v = \text{unit} \cdot (x - c_t)^{m_{crit}(c)+1}, \quad (x, t) \text{ near } (c_t, t), \quad (10.1)$$

Then $\frac{\partial}{\partial t}$ lifts to a unique real analytic vector field in $\mathbb{R} \times I_{\mathbb{R}}$ which generates $\mathcal{D}$.

For the case $\mathbb{K} = \mathbb{C}$, we still have d , which is (merely) real analytic, and also (10.1), where c_t is holomorphic in t . Thus $\frac{\partial}{\partial t}$ admits a *local* real analytic lifting. By Cartan's Theorem B, a real analytic *global* lifting exists. This completes the proof. $\square$

11. APPENDIX 2. TREE MODELS

The *tree-model* ([13]) is best explained by an example. Take $\phi(\xi) = (\xi^2 - y^3)^2 - 4\xi y^5$. The Puiseux roots are

$$\zeta_i(y) = \pm y^{3/2} \pm y^{7/4} + \cdots, \quad 1 \leq i \leq 4.$$

The tree-model is shown in Fig. 3. Tracing upward from the tree root to a tip along solid line segments amounts to identifying a Puiseux root.



There are three bars, of height $3/2$, $7/4$, $7/4$ respectively. Each is indicated by a horizontal line segment (whence the name); the associated polynomial has *at least two* distinct roots.

At these heights, the ζ_i 's split away from each other. *Bars without this property are not indicated*; they correspond to the *vertex edges*, Fig. 4.

The three critical points (polars) γ_j are indicated by dashed lines; their positions relative to the ζ_i are justified by Theorem 5.3, exposing the contact orders.

The Newton Polygon $\mathcal{NP}(\phi, \zeta_i)$ is shown in Fig. 4. The dotted segments e_1, e_2, e_3 indicate vertex edges, where

$$h(e_1) < 3/2 < h(e_2) < 7/4 < h(e_3) < \infty.$$

FIGURE 3. Tree Model $\mathbb{M}_{1,\phi}$ FIGURE 4. $\mathcal{NP}(\phi, \zeta_i)$

REFERENCES

1. Abderrahmane J, Ould M., *Newton Polygon and Trivialisation of Families*, J. Math. Soc. Japan, 54, 513-550 (2002).
2. K. Bekka and S. Koike, *The Kuo condition, an inequality of Thom's type and (C)-regularity*, Topology, **37** (1998), 45-62.
3. E. Bierstone and P. D. Milman, *Arc-analytic functions*, Invent. math., **101** (1990), 411-424.
4. E. Bierstone, P. Milman *Semianalytic and subanalytic sets*, Inst. Hautes tudes Sci. Publ. Math. No. 67, 1988, 5-42.
5. J. Bochnak, S. Lojasiewicz *A Converse of the Kuiper-Kuo Theorem*, in Proc. Liverpool Singularities Symposium, I, (ed. C.T.C. Wall), 1971, Lecture Notes Math., 209, Springer-Verlag.
6. E. Brieskorn, H. Knörrer, *Plane Algebraic Curves*, Birkhäuser Verlag, 1986.
7. Charles Favre, Mattias Jonsson, *Valuative analysis of planar plurisubharmonic functions*, Invent. Math. 162 (2005), No. 2, pp 271-311.

8. T. Fukui and L. Paunescu, *Modified Analytic Trivialization for Weighted Homogeneous Function Germs*, J. Math. Soc. Japan, vol. 52, No. 2, (2000), pp 433–446.
9. H. Hancock, *Theory of Maxima and Minima*, Dover Publications, Inc. New York 1960.
10. Theo de Jong, Gerhard Pfister, *Local analytic geometry. Basic theory and applications*, Advanced Lectures in Mathematics. Friedr. Vieweg & Sohn, Braunschweig, 2000.
11. S. Koike, *Notes on C^0 -sufficiency of quasijets*, J. Math. Soc. Japan, **42** (1990), 265–275.
12. T.-C. Kuo, *On C^0 -sufficiency of jets of potential function*, Topology 8, p.157–171 (1969).
13. T.-C. Kuo and Y.C. Lu, *On analytic function germs of two complex variables*, Topology, 15 (1977), 299–310.
14. T.-C. Kuo and A. Parusiński, *Newton Polygon relative to an arc*, in Real and Complex Singularities (São Carlos, 1998), Chapman & Hall Res. Notes Math., **412**, 2000, 76–93.
15. T.-C. Kuo and L. Paunescu, *Equisingularity in R^2 as Morse stability in infinitesimal calculus*, (Communicated by H. Hironaka) Proc. Japan Acad. Ser. A, Math. Sci. Volume **81**, Number 6 (2005), 115–120.
16. J. Milnor, *Morse Theory*, Annals of Math. Studies. **51**, Princeton Univ. Press 1983.
17. J. Milnor, *Singular Points of Complex Hypersurfaces*, Annals of Math. Studies. **61**, Princeton Univ. Press 1968.
18. L. Paunescu, *V-sufficiency from the weighted point of view*, J. Math. Soc. Japan, vol. 46, No. 2, (1994), pp 345–354.
19. L. Paunescu, *A Weighted Version of the Kuiper-Kuo-Bochnak-Lojasiewicz Theorem*, J. of Algebraic Geometry, volume 2, number 1, January (1993), pp 69–79.
20. F. Pham, *Deformations equisingularités des idéaux jacobiens de courbes planes*, in Proc. Liverpool Singularities Sym., II, (ed. C.T.C. Wall), 1971, 218–233, Lecture Notes Math., 209, Springer-Verlag.
21. van der Waerden, *Einführung in die Algebraische Geometrie*, Zweite Auflage. Die Grundlehren der Mathematischen Wissenschaften, Band 51. Springer-Verlag, Berlin-New York, 1973.
22. R. J. Walker, *Algebraic Curves*, Springer-Verlag, 1972.
23. H. Whitney, *Complex Analytic Varieties*, Addison-Wesley, Reading, Mass., 1972.
24. O. Zariski, *Algebraic Surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 61. Springer-Verlag, New York-Heidelberg, 1971.
25. O. Zariski, *Studies in equisingularity I*, Amer J. Math. 2, **87**, (1965), 507–536.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, SYDNEY, NSW, 2006, AUSTRALIA

E-mail address: tck@maths.usyd.edu.au; laurent@maths.usyd.edu.au